

Fish Oils-Volume I #104 12/5/73

FISH OILS VOL I #104

M23

VOLUME 1

GRAS MONOGRAPH SERIES

FISH OILS

prepared for
THE FOOD AND DRUG ADMINISTRATION
DEPARTMENT OF HEALTH, EDUCATION
AND WELFARE

DECEMBER 5, 1973

This publication was prepared under Contract Number FDA 72-100
with the Public Health Service, Food and Drug Administration,
Department of Health, Education, and Welfare

prepared by
Tracor Jitco, Inc.

Table of Contents

Summary

CHEMICAL INFORMATION

	Page
I. Nomenclature	1
II. Empirical Formula	2
III. Structural Formula	12
IV. Molecular Weight	12
V. Specifications	12
VI. Description	13
VII. Analytical Methods	17
VIII. Occurrence and Levels Found in Plants and Animals	17

BIOLOGICAL DATA

I. Acute Toxicity	20
II. Short Term Studies	20
III. Long Term Studies	27
IV. Special Studies	27

BIOCHEMICAL ASPECTS

I. Breakdown	29
II. Absorption-Distribution	31
III. Metabolism and Excretion	33
IV. Effects on Enzymes and Other Biochemical Parameters	36
V. Drug Interaction	37
VI. Consumer Exposure Information	38

Bibliography

Journal article copies and translations

SUMMARY

Description and Specifications

For present purposes fish oil may be defined as the oil expressed primarily from menhaden, and secondarily from herring or tuna (839). Outside the United States, herring is currently the primary source (828). The California sardine (pilchard) was finally overfished in 1951 (839), the Peruvian anchovy in 1972, probably (386). Whale oil can no longer be used legally in the United States.

Fish oils are especially rich in long chain polyunsaturated fatty acids (PUFA), with more than C_{18} and with four or more double bonds (049,305,836). About 2% of their contents have essential fatty acid (EFA) activity (730). They also contain small amounts of volatile short chain acids (176), phospholipids, ether-linked compounds, hydrocarbons, wax esters and sterols, such as cholesterol and "vitamin" D (558,836,501,496).

The purpose of hydrogenation is to elaborate a stable, palatable product of suitable plasticity (028). Hydrogenation is a variable process that saturates some but not all of the double bonds; the aim is to partly-saturate polyene chains but not monoene (028). Therefore, hydrogenated fish oils are never more than partly hydrogenated. The proportion of double bonds left unsaturated is reflected by the iodine number. But the fatty acid (FA) composition is altered, in most cases not very predictably (028,324,502,635,864).

Manufacturers patent their particular hydrogenation procedures to protect products with special handling characteristics (621,624,625,649). Thus, hydrogenated fish oils include a wide range of chemically different products that vary according to source and subsequent treatment (864).

The two important food uses of hydrogenated fish oils are in margarines and shortenings. Currently in the United States, these oils are used only in non-food industrial products for example, in paints and lubricants. This situation may change, however (828,836). (See below, Consumer Exposure.)

At present there is no standard of identity for hydrogenated fish oils that may be used in the United States' food supply (188,053,898,899,828). The nearest approach is the FAO/WHO Codex Alimentarius Recommended International General Standard for Edible Fats and Oils Not Covered by Individual Codex Standards (181). This sets maxima for acid and peroxide values, stabilizers, and other additive contents, and for contaminants such as lead, copper, arsenic, iron, and soap.

Part of the trouble has been a lack of usable routine analyses for the major components of hydrogenated fish oils. Very recently, development of sophisticated fractionation procedures and variants of gas-liquid chromatography have begun to overcome this lack (560,561,564, 718,380,028,835). Also fish oils have been regarded as

auxiliary sources of fats that are major nutrients, rather than as separately identifiable additives. From this viewpoint, hydrogenation introduces an uncontrolled and unwanted variable into research studies.

Consequently, the literature contains a great deal of information on fish oils but relatively little on hydrogenated fish oils (901,622,502). This monograph necessarily draws on the former for supportive data.

Acute Toxicity

Only one series of studies on the acute toxicity of fish oils was found. Between 1954 and 1962, Matsuo, in Japan, searched the degradation products of cuttlefish oil and found a cyclic monomer that was acutely toxic to rats and rabbits. He concluded that this monomer was an end product of autoxidation and peroxidation of many edible oils. The starting material was not hydrogenated (585,586, 597,598,599).

Short Term

Studies

No studies on hydrogenated fish oils in man were found. No animal studies were found on hydrogenated fish oils used to supplement normal diets. Most studies that were identified are on fresh fish oils, on fractions of fish oils, on FA that occur in fish and other oils, or on derivatives and degradation products where the starting material happened to be a fish oil. In some cases defatted or semipurified diets were supplemented with

large amounts of the experimental substances, or of a hydrogenated fish oil.

Reports summarized in this section of the monograph include those of Haven (345), Dam (204), Blaxter et al. (111,112,113), Dam (204), Ershoff (251), Andrews et al. (058), Gershoff and Norkin (286), Rasheed et al. (739), Carpenter et al. (163), Roubal and Tappel (760), Stansby (832), Njaa et al. (668), and Grover (303).

From these studies emerged three major conclusions, applying to hydrogenated fish oils and all other dietary fats and oils:

1. Toxic effects reflect mainly deficiencies resulting from imbalances in the total experimental diets. In many studies the deficiency was of EFA activity (204,111,007,836).
2. Increased intakes of PUFA increase the requirements for vitamin E (112), largely but not wholly for its antioxidant properties (112,739). Calculations by Harris and Embree (333) suggested that in 1960 the population of the United States (consuming no fish oil) received only marginally sufficient vitamin E. However, this has been denied recently by the National Academy of Sciences (657). According to Hannewijk (324), carefully refined and hydrogenated fish oils contain adequate antioxidant activity. Many of the studies that were reviewed, however, reported toxic symptoms of vitamin E deficiency (202,286).

3. If oil putrefies by autoxidation, peroxides form, and if these are not destroyed in the intestine, results can be toxic (058,739). For example, free radicals arise that can damage proteins and amino acids (204,760). The more highly unsaturated the PUFA, the more readily the oil autoxidizes (832). According to Stansby (832) these are "all only potential effects" that can be "controlled by the use to which the oil is put."

Theoretically, peroxides can give rise to epoxides, which can be carcinogenic (303). But no studies were found to suggest that this might occur in vitro or in vivo with hydrogenated fish oils. On the contrary, Haven (345) reported that induced cancer in rats grew more slowly when the dietary fat was an extremely hydrogenated fish oil than when it was a highly unsaturated vegetable oil.

Long Term Studies

No relevant long-term studies were found in the literature. The controversial question of the general relationships of saturated/unsaturated fats, cholesterol, and atherogenesis was considered to be outside the scope of a monograph restricted to hydrogenated fish oils.

Special Studies

Evidence from the U.S. National Marine Fisheries Service (848,849) is reviewed to the effect that in some areas sea fish contain unacceptable levels of DDT and related pesticides, attributable to agricultural run-off. Stout and her colleagues (849) have established that these residues are substantially eliminated by the normal refining

and hydrogenation of fish oils. However, problems with the detection and elimination of PCB's remain (094).

Breakdown and

Metabolism

Oxidation results in "off" flavors, and hydrogenation generates "hardening" flavors that are removed by further refining. The latter arise from small amounts of free or bound aldehydes (174), especially 6-trans-nonenal, which can sometimes be detected at a dilution of 0.3 ppb (620, 458,324). Off or hardening flavors can return during storage, and such oils become unacceptable to the taste before they become toxic (828). Stansby (828) notes that hydrogenated menhaden oil is no exception but, being only moderately unsaturated, should cause no special problems.

No reports were found to suggest that the digestion, absorption, or distribution of hydrogenated fish oils might differ from those of other dietary fats. However, a paper by McCay and Paul (607) suggests that hydrogenation may diminish absorption of any oils and thus increase the amounts excreted in the feces. In general, the FA distribution in the tissues tends to reflect that in the diet, taking into account that certain FA are readily metabolized or liable to inhibit the metabolism of other FA. This was shown in rats fed hydrogenated herring oil (090), in turkeys fed anchovy oil (660), and in Japanese fishermen with high intakes of fish oils (388).

Although EPA activity of fish oils (unhydrogenated) is equivalent to that of corn oil, a common source of EPA (730), much of this activity seems to be lost on hydrogenation (007); the reasons are not clear, however, and further research has been recommended (007). Losses of vitamins, including vitamin E, through hydrogenation also appear to require further study (913); van der Steur (913) has pointed to risks if they are over-replaced.

The literature contains many reports of metabolic effects of fish oils, and of the presence or lack of FA that are found in fish oils--see especially a cholesterol-lowering effect reviewed by Stansby (836)--but there is no estimate of the relevance of these reports to hydrogenated fish oils. This monograph does not survey the vast field of knowledge of FA metabolism and its interactions with other nutrients and metabolic pathways.

Drug Interactions No reports have been found.

Consumer Exposure Until the disappearance of the California sardine in 1951, large amounts of hydrogenated fish oils were made into margarine (839). Then the use of fish oils ceased; in 1955, when the present FDA standard of identity for oleo-margarine (067) was set up, nobody asked for fish oils to be included (839). Today, therefore, hydrogenated fish oils cannot legally be included in margarine in the United States, although they are used extensively abroad (839).

CHEMICAL INFORMATION

I. Nomenclature

- A. Common Names: Fish Oil (Hydrogenated); Hydrogenated Fish Oil.
- B. Chemical Names: None for the entire product.
- C. Trade Names: Either the common names are used or the species of fish that yielded the oil in the sample referred to. Examples are: Menhaden Oil, Herring Oil, Tuna Oil, etc. qualified by the adjective "hydrogenated," or "partly hydrogenated", or even "partially (sic) hydrogenated".

Currently, the menhaden is the predominant species of fish used as a source of fish oil (hydrogenated) in the United States. Four principal species of menhaden have been identified in the areas of the U.S. fisheries, two in the Atlantic Ocean and two in the Gulf of Mexico. The two commonest are Brevoortia tyrannus in the Atlantic, and B. patronus in the Gulf of Mexico (839). No references, however, were found in which a more precise term was used than "menhaden oil".

- D. Chemical Abstracts Services Unique Registry Number: MX 8016 14 6.

NOTE: Relatively few reports on hydrogenated or partly hydrogenated fish oils were found, compared to the number on unhydrogenated fish oils. Some of the latter reports are therefore included as supporting data.

II. Empirical Formula

The composition of fish oil varies greatly according to its source. Stansby (836) points to "vast differences in composition of oils from different species of fish or even from oils of the same species of fish taken from different geographical areas," and also from different parts of the same fish. Major species of fish used as oil sources have tended to vanish, because of overfishing. One example was the Hokkaido herring. California sardine oil was last available in 1951 (828), and 1972 may have seen the end of the Peruvian anchovy as a major source of oil (386). Currently, the menhaden is the major source of fish oil in the United States, but production declined between 1960 and 1967 (839). Elsewhere the herring is the major oil source. Tuna is a less important source.

Fish liver oils, usually sold as carriers of vitamins A and D in amounts that are now categorized as "drugs" or "diet supplements", are outside the scope of this monograph. Whales are now "endangered species" whose oils cannot legally be marketed in the United States.

Table 1 shows the fatty acid (FA) compositions of menhaden and herring oils according to Altman and Dittmer (049). Compositions listed by Gruger (305) and by Stansby (836) differ considerably, but all authors have emphasized variability. Some of these fatty acids are described more fully in Tables 2 and 3 (from Altman and Dittmer 049). Fish oils are especially rich in long chain polyunsaturated fatty acids (PUFA).

The essential fatty acid (EFA) properties of the three fish oils most important to the fish oil production of the United States (menhaden, herring, and tuna oils) were investigated by Privett et al. (730).

Table 1

Typical Fatty-Acid Composition of Two Fish Oils*

FASEB Serial No.	Common name of fatty acid	Concentration g/100 g total fatty acids
<u>Menhaden Oil</u>		
(74)	Myristic acid	5.9
(75)	Palmitic acid	10.3
(76)	Stearic acid	0.6
(77)	Arachidic acid	0.6
(78)	Behenic acid	0.8
(79)	Palmitoleic acid	15.5
(80)	Linolenic acid	29.6
(81)	Eicosapolyenoic acid	19.0
(82)	Docosapolyenoic acid	11.7
<u>Herring Oil</u>		
(67)	Myristic acid	7.3
(68)	Palmitic acid	13.0
(69)	Stearic acid	Trace
(70)	Palmitoleic acid	4.9
(71)	Linolenic acid	20.7
(72)	Eicosapolyenoic acid	30.1
(73)	Docosapolyenoic acid	23.2

* Adapted from R.P. Geyer (Table 43) in Altman and Dittmer (1949).

Table 2

Fatty Acids Found in Fish Oils*

FASEB Serial No.	Identity	Common name	Empirical formula	Sources
<u>Saturated</u>				
(14)	Tetradecanoic	Myristic	$C_{14}H_{28}O_2$	see Table 1
(16)	Hexadecanoic	Palmitic	$C_{16}H_{32}O_2$	see Table 1
(18)	Octadecanoic	Stearic	$C_{18}H_{36}O_2$	see Table 1
(20)	Eicosanoic	Arachidic	$C_{20}H_{40}O_2$	see Table 1
(22)	Docosanoic	Behenic	$C_{22}H_{44}O_2$	see Table 1
<u>Unsaturated - monoethenoic</u>				
(95)	<u>cis</u> -9-Hexadecenoic	Palmitoleic	$C_{16}H_{30}O_2$	see Table 1
(124)	<u>cis</u> -11-Eicosenoic	<u>cis</u> -Gondoic	$C_{20}H_{38}O_2$	Menhaden
(125)	<u>trans</u> -11-Eicosenoic	<u>trans</u> -Gondoic	$C_{20}H_{38}O_2$	Menhaden
(128)	<u>cis</u> -13-Docosenoic	<u>cis</u> -Erucic	$C_{22}H_{42}O_2$	Herring (305)
<u>Unsaturated - polyenoic</u>				
(169)	11,14-Eicosadienoic		$C_{20}H_{36}O_2$	Herring, Menhaden

Fatty Acids Found in Fish Oils (Cont'd)

(170)	5,13-Docosadienoic		$C_{22}H_{40}O_2$	Fish oils
(172)	17,20-Hexacosadienoic		$C_{26}H_{48}O_2$	Fish oils
(175)	6,9,12-Hexadecatrienoic		$C_{16}H_{26}O_2$	Herring, Menhaden
(177)	7,10,13-Hexadecatrienoic		$C_{16}H_{26}O_2$	Menhaden
(180)**	<u>cis-6,cis-9,cis-15</u> -Octadecatrienoic	γ -Linolenic	$C_{18}H_{30}O_2$	see Table 1
(191)**	<u>cis-9,cis-12,cis-15</u> -Octadecatrienoic	α -Linolenic	$C_{18}H_{30}O_2$	see Table 1
(196)	5,8,11-Eicosatrienoic		$C_{20}H_{34}O_2$	Fish oils
(197)	8,11,14-Eicosatrienoic		$C_{20}H_{34}O_2$	Fish oils
(205)	6,9,12,15-Hexadecatetraenoic		$C_{16}H_{24}O_2$	Herring, Menhaden
(214)	6,10,14,18-Eicosatetraenoic		$C_{20}H_{32}O_2$	Fish oils
(222)	4,8,12,15,19-Docosapentaenoic	Clupadonic	$C_{22}H_{34}O_2$	Fish oils
(223)	7,10,13,16,19-Docosapentaenoic		$C_{22}H_{34}O_2$	Herring

* Extracted from K.S. Markley (Table 42) in Altman and Dittmer (1949), except where otherwise stated.

** Linolenic acid is stated to have two isomeric forms (α and γ) with indistinguishable properties.

Table 3

Some Chemical Characteristics of Fatty Acids Found in Fish Oils*

FASEB Serial No.	Iodine No.**	Molecular weight	Melting point °C	Boiling point °C (at mm Hg)	Specific gravity (at °C)***	Refractive index**** (at °C)	Soluble in*****
(14)	0	228.4	53.9	250(100)	0.8622(54)	1.4273(70)	acet,alc,eth,pet.eth.
(16)	0	256.5	63.1	268(100)	0.8487(70)	1.4309(70)	acet,halc,eth,pet.eth.
(18)	0	284.5	69.6	213(5)	0.8390(80)	1.4337(70)	acet,halc,eth,pet.eth.
(20)	0	312.5	76.5-77.0	201(1)	0.8240(100)	1.4250(100)	bz,ch1,eth,pet.eth.
(22)	0	340.6	81.5	306(60)	0.8221(100)	1.4270(100)	sl.sol:alc,eth.
(95)	99.77	254.40	-0.5 to +0.5	140-141(5)	-	-	-
(124)	81.74	310.50	23-24 50	267(15)	-	-	alc,me.alc.
(125)	81.74	310.50	52-53	-	-	-	-
(128)	74.97	338.56	34.7	281(30)	0.8352(70)	1.4444(70)	eth,me.alc.
(169)	164.55	308.49	-	-	-	-	acet,eth,pet.eth.
(170)	150.83	336.54	-	-	-	-	-
(172)	129.28	392.64	61	-	-	-	eth,pet.eth.
(175)	304.12	250.37	-	-	-	-	alc,eth.
(177)	302.12	250.37	-	-	-	-	acet,alc,eth,pent.
(180)	273.51	278.40	-11.3 to -11	125(0.05)	0.9164(20)	1.4800(20)	acet, me.alc,eth,pet.eth.

Table 3 - Cont'd.

(191)	273.51	278.40	-11.3 to -11	125(0.05)	0.9164(20)	1.4678(50)	acet,alc,eth,pet.eth.
(196)	248.48	306.45	-	-	-	-	CS ₂ ,me.alc,hept.
(197)	248.48	306.45	-	-	-	-	CS ₂ ,me.alc,hept.
(205)	407.98	248.45	-	-	-	1.4870(29)	CS ₂ ,acet,alc,eth,pent.
(214)	333.48	304.44	-	-	0.9263(20)	1.4935(20)	acet,me.alc,eth,pet.eth.
(222)	384.0	302.49	-78	207-212(2)	0.9356(20)	1.5014(20)	acet,eth,pet.eth.
(223)	384.0	332.49	-	-	-	-	bz,chl,me.alc,pet.eth.

* Identified in Table 2, cf. FASEB serials Numbers

** Grams of I absorbed by 100 g of acid.

*** Referred to water at 4°C.

**** At the D-line of Na (589 nm).

***** sl = slightly; sol = soluble; acet = acetone; alc = alcohol (ethyl unless otherwise stated); h = hot;
me = methyl; eth = diethyl ether; pet = petroleum; h = heptane; pent = pentane; bz = benzene; chl = chloroform.
Gaps mean no data reported in Table 42 of Altman and Dittmer (1949).

using two groups of 50 rats. They found that about 2% of each of these oils could be categorized as EFA using the dermal symptoms of EFA deficiency as the criterion.

Fish oils also contain volatile short-chain acids. Chipault and McMeans (176) collected 25 ml of volatile compounds during distillation of 5 gallons of menhaden oil. Paper chromatography indicated that these included formic (or acetic), acrylic, propionic, crotonic, butyric, and valeric acids; not all of the compounds were identified.

Despite their variability, fish oils as a class are distinguished by their high contents of fatty acids with chains longer than 18 carbons, and with four or more double bonds. Most fish oils are also rich in lipids other than FA and triglycerides: these include phospholipids, ether-linked compounds, hydrocarbons, wax esters, and sterols. In reviewing the compositions of fish oils Malins (558) and Stansby (836) remarked on the incompleteness of current knowledge. In general, lecithin and cephalin together account for about 75% of the phospholipids; most of the ether-linked compounds are found as diacyl glyceryl ethers; squalene is said to be the principal hydrocarbon and sterols comprise mainly cholesterol and related substances including "vitamin" D. Table 4 illustrates the FA composition of lecithins in menhaden, salmon and tuna oil.

Lambertsen and Holman (501), partly characterized the hydrocarbons of herring oil by gas chromatography after various separation procedures. Among the straight-chain paraffins the odd carbon numbers were found to predominate, especially 27.0; among the branched-chain hydrocarbons, pristane (16.8) was 70%, and squalene (26.3) only 13% (Table 5).

Table 4

Specific Distribution of the Principal Fatty Acids from the Lecithins of Fish (622)

Fatty Acids	Weight Per Cent of Specific Fatty Acid Isolated From								
	Salmon			Menhaden			Tuna		
	Total	α'	β	Total	α'	β	Total	α'	β
14:0	0.7	1.9	0.6	0.5	Tr	1.3	0.7	Tr	1.0
14:1	—	—	—	Tr	Tr	Tr	Tr	Tr	1.8
16:0	22	37	10	45	78	6.2	19	36	5.0
16:1	8.0	1.4	3.8	Tr	0	Tr	2.1	Tr	2.3
18:0	2.0	2.7	Tr	3.2	5.9	0.9	5.0	7.9	1.0
18:1	6.7	8.2	23	7.4	2.3	13	7.6	6.4	16
18:2	Tr	Tr	Tr	1.0	Tr	2.4	0.7	Tr	2.2
18:3	Tr	Tr	Tr	Tr	Tr	Tr	0.8	Tr	1.1
20:4	1.0	0.6	0.3	2.2	Tr	0.9	2.7	1.3	5.0
20:5	12	14	17	13	1.7	29	7.1	8.4	15
22:1	0	0	0	0	0	0	0	0	0
22:4	Tr	Tr	Tr	Tr	Tr	Tr	Tr	0	0.5
22:5	Tr	Tr	Tr	Tr	Tr	Tr	Tr	Tr	Tr
22:6	43	33	46	28	12.5	42	55	39	48

From Menzel and Oleott (1961).

Table 5

Hydrocarbons Occurring in Herring Oil as Determined by Gas-Liquid Chromatography (501)

Straight-Chain Components		Branched-Chain Components	
Carbon Number	% Area	Carbon Number	% Area
14.0	0.3	15.4	3.5
15.0	7.5		
16.0	0.6	16.8 (pristane)	70.0
17.0	9.0		
17.6	2.2	17.9	5.0
18.0	1.7		
19.0	10.5	19.0	0.5
20.0	0.3		
21.0	6.0	19.9	1.0
22.0	0.3		
23.0	0.8	21 ¹	3.5
24.0	0.3		
25.0	9.5	22.2	3.5
26.0	Trace		
27.0	33.0	26.3 (squalene)	13.0
28.0	Trace		
29.0	9.0		
30.0	Trace		
31.0	~7		
32.0	(Trace)		
33.0	~1		

¹ Flat and broad peak.

From Lambertsen and Holman (1963).

Kritchevsky et al. (496) studied the nonsaponifiable fraction of various sea foods for sterol content, and found that in three teleosts (haddock, pollock, and salmon) cholesterol accounted for more than 90%.

The effects of catalytic hydrogenation on these lipids have received relatively little study. Lambertsen et al. (503) took five samples of industrially hydrogenated marine oils, fractionated them by thin-layer chromatography, and analyzed the fractions by gas-liquid chromatography. "The general fatty acid composition was 30-45% saturated acids, 45-55% monoenoic acids, and 10-20% polyenoic acids. A substantial part of the polyenoic acids were trienoic acids, the rest dienoic. The presence of tetraenoic acids could be ascertained in the low-melting samples (mp 30-32°C)."

Ackman et al. (028) stated: "The partial hydrogenation of fats and oils to produce a stable, palatable product of suitable plasticity results in the formation of new, structurally altered fatty acids which in most cases differ chemically from familiar natural products.' The chemical aim of hydrogenation is selectively to saturate the polyene chains, but not the monoene.

Two other effects of hydrogenation have been reviewed by Hannewijk (324)--migration of double bonds, and cis-trans isomerization. These effects have been studied lately in monoene FA by Lambertsen et al. (502) and by Ackman et al. (028). No reports were found of studies on polyene FA.

Thus, both the original composition of fish oil and the effects of part-hydrogenation are variable and somewhat unpredictable. Therefore, the composition of Fish Oil (Hydrogenated) cannot be described within

narrow limits of components and their quantitative distribution. Large variations among samples can be expected. Nevertheless, the hydrogenation of fish oils does not make them resemble other hydrogenated oils. This was demonstrated long ago by an elegant series of cooling curves (900). Nor do hydrogenated oils from different fish species come to resemble each other chemically (864).

III. Structural formula

No single structure exists. Examples of the structures of some components of fish oils are given in Figure 1.

IV. Molecular weight

Data for the unhydrogenated FA components of fish oils are given in Table 3. Molecular weights of hydrogenated fish oils vary according to composition and degrees of hydrogenation.

V. Specifications

Fish Oil (Hydrogenated) is not mentioned in the Food Chemicals Codex, 2nd Edition, 1972 (188), the National Formulary, 13th Edition, 1970 (053), the U.S. Pharmacopoeia, 18th Edition, 1960 (898), or Synthetic Organic Chemicals, U.S. Production and Sales (U.S. Tariff Commission) (899). Nor are fish oils mentioned (unqualified by the word "hydrogenated").

According to Stansby (828), Codex Alimentarius standards have been recommended for fish oils. However, the current FAO Books in Print (267) contains no individual Codex Alimentarius standard for fish oils. Therefore, fish oil (hydrogenated) will be covered by FAO publication

No. CAC-RS 19-1969, General Standard for Edible Fats and Oils Not Covered by Individual Codex Standards (181).

Specifications relevant to the identity and toxicity of hydrogenated fish oils found in CAC-RS 19-1969 include the following:

3.5 Peroxide Value not more than 10 milliequivalents of peroxide oxygen/kg fat or oil

5.5 Copper (Cu): Non-virgin oil: 0.1 mg/kg

5.7 Arsenic (As): 0.1 mg/kg

7.1.2 Where an oil has been subject to any process of esterification or to processing which alters its fatty acid composition or its consistency the specific name of the oil shall not be used unless qualified to indicate the nature of the process.

VI. Description

Table 3 gives some physical properties of fatty acids of unhydrogenated fish oils; there are many gaps in the data. Table 6 similarly illustrates some properties of entire unhydrogenated fish and other oils. Hydrogenation alters these properties and also the flavor of fish oils. Figure 2, taken from Chang (172), summarizes the processing of fish oils.

Hannewijk (324) has reviewed the hydrogenation process. Oils are hardened to an over-all melting point (mp) of 35-37°C, monitored by sampling. A crude oil with more than 0.1% of FFA must be bleached and neutralized before hydrogenation. Other problems include high levels of bound aldehydes; however, most impurities are removed by normal refining. Added H₂ should be free from S and CO. If oil becomes hotter

Table 6

Some Physical and Chemical Constants of Fish Oils and Other Oils

Type of oil	$n_D^{65}=1.4^*$ limits mean	Iodine value limits mean	Slip point(°C) limits mean	Saponification value limits mean
<u>Stansby (833)</u>				
Herring	573-602 565	123-148 134	9-17 14	181-187 184
Menhaden	590-623 608	150-165 159	22-28 24	192-199 196
<u>Fineberg and Johansen (263)</u>				
Herring		120-150		185-195
Menhaden, Atlantic		150-185		185-190
Menhaden Gulf		145-170		**
Soybean		120-141		189-195
Rapeseed		97-108		170-180

* "A relationship exists between iodine value and refractive index such that $I.V. = (n_D^{65} - 1.4429) \times 8560$." (833).

** Reliable data said to be unavailable (263).




Figure 2. Flow Sheet of Fish-Oil Processing. From Chang (172).

than about 200°C, aromatic FA (bound to glycerol) tend to be formed; however, high temperatures improve the selective hydrogenation of higher PUFA. A solid, preferably Ni-containing, catalyst of known pore size is used. Its contamination with S enhances cis-trans isomerization resulting in a fat that is firmer at room temperatures; however, S contamination accumulates, and must be watched. The kinetics of hydrogenation have been studied in detail by Miyake et al. (635). Koritala (491) has described Cu-containing catalysts, and Beal et al. (688) give a method for removing Cu residues from hydrogenated oils.

So the hardening of oils by hydrogenation results from two sorts of chemical alterations--saturation of double bonds, and cis-trans isomerization. The relative amounts of these alterations and their control, are the subject of patents by which manufacturers seek to guard the individuality of their branded products (621,624,625,649). So, also, hydrogenated oils and fats are mixtures of lipids with very different mp and expansion spectra, and the latter are described as "steep" or "flat".

The hydrogenated fats are treated with weak alkali to remove traces of FFA and the catalyst, and are then bleached and deodorized. According to Hannewijk the finished oils or fats should be stable for about three months, which is longer than the usual shelf-life of margarine in Western Europe (for American practice, see Consumer Exposure). If the oils are processed carefully, antioxidants are not needed; if not, they are of little avail. This was also the opinion of Lea (512),

"Hardening flavors" resulting from hydrogenation are attributed to the presence of free ketones and especially free aldehydes, for which

taste thresholds can be very low, e.g., 0.0003 ppm for 6-trans-nonenal. Much higher levels of some aldehydes can develop within hours in daylight and within about two months in the dark. Hardening flavor also varies with the mp spectrum and the degree to which PUFA are saturated. Furthermore, human taste thresholds vary enormously, see review by Carr et al. (164). Hannewijk states that hardening flavors can almost always be perceived in herring oils of mp 30°C or less, are "largely removed" at mp 33°C, but are occasionally present even at mp 35°C. Menhaden oil, however, must be hardened to mp 40°C or more to eliminate hardening flavors.

All authors emphasize that most of the keeping qualities of hydrogenated fish oils depend on careful refinement before hydrogenation.

VII. Analytical methods

The CAC General Standard for Edible Fats and Oils (181) lists acceptable methods for determining acid value; peroxide value; matter volatile at 105°C; insoluble impurities; soap content; and contents of iron, copper, lead, and arsenic. No specific CAC Standard has yet been issued for fish oils or hydrogenated fish oils.

Malins and Mangold (560) and Mangold (564) described the analysis of complex lipid mixtures by methods centering on thin-layer chromatography (TLC). In a later paper, Malins and Wekell (561) described the classification of lipids by adsorption chromatography, using columns or TLC and a variety of adsorbents. Many other fractionation procedures are used for determining useful components of fish oils; for example, Pohle et al. (718) reported measurement of solids by nuclear magnetic resonance.

Houle (380) has determined the FA composition of fish oils by gas-liquid chromatography. Ackman et al. (028) used open tubular gas-liquid chromatography to analyze a hydrogenated herring oil of mp 32.5°C, iodine number 76, 30% saturated FA, and 66% monounsaturated FA. They found that they could determine the cis and trans isomers.

Stansby (835) reviewed procedures for analyzing complete fish oils for saturation (iodine number), FFA, moisture, color, and impurities, by standard procedures listed by the Association of Official Agricultural Chemists (070) and in the Official and Tentative Methods of the American Oil Chemists Society (052).

Flavors, however, cannot be determined adequately without using a human sensory evaluation panel. Such a panel cannot render determinations or exact measurements (164).

Banks (082) reviewed deteriorative changes in fish oils, including the slow formation of FFA by hydrolysis in the presence of alkali or of lipases (enhancing autoxidation), and the more important autoxidation that can lead to formation of hydroperoxides and thence epoxides (see Section B. II). The standard test for determining oxidation products of fish oils at present is the 2-thiobarbituric acid test (222).

Impurities arising from the environment or from processing of fish oils include copper, arsenic, and bromine. Analyses more recent than, or not included in, the CAC General Standard (181), that have been found in the literature, are as follows:

Deck and Kaiser (225) developed a colorimetric analysis of edible oils for copper content. Lunde (541) described a neutron activation

analysis for arsenic and bromine after fractionation of oils on silica-gel columns.

All of the above methods are described for fish oils in general. The following methods have been described for hydrogenated fish oils:

Ueno (901) determined thiocyanogen value as a measure of the degree of hydrogenation. Sardine oil showed a rapid decrease of thiocyanogen value at the stage of hydrogenation where the iodine number fell to about 70-80.

Menzel and Olcott (622) described an enzymatic analysis for positional distribution of FA in the phosphatidyl choline (lecithin) fractions from various animals and fish including tuna and menhaden.

Lambertsen et al. (602) lately developed a fractionation procedure and gas-liquid chromatographic analysis for monoene FA isomers in hydrogenated fish oils.

VIII. Occurrence and Levels Found in Plants and Animals

- A. Plants. Not applicable. However, it is well known that some, if not all, fatty acids can move up the food chain unaltered.
- B. Animals. See Sections II and VI above.
- C. Synthetics. Not applicable.
- D. Natural Inorganic Sources. Not applicable.

BIOLOGICAL DATA

I. Acute Toxicity

Matsuo studied the toxicity of fish oils for many years. His first reported finding was that rats fed diets containing 5% of ethyl esters of highly unsaturated FA from cuttlefish oil thrived, but the oxidized forms of these esters were toxic (586). The degree of toxicity was found to be directly proportional to the peroxide content which could result from autoxidation of the oil (585). "Profound damage" to the mucous membranes of stomach and intestines of rabbits, and liver damage, resulted from oral administration of 0.8 ml per day of the autoxidized esters, and topical application of these or non-oxidized esters to the back skin of rats had like effects (598). Rats were similarly affected by peroxides formed in cod liver oil by autoxidation (597). After various experiments involving heat-polymerization of cuttlefish oil and of ethyl linolenate, both of which gave rise to cyclic esters that proved to be acutely toxic to rats, Matsuo purified a cyclic monomer of mw 282. This proved rapidly fatal to rats when fed as 20% of a basal diet, and Matsuo concluded that this monomer was the toxic endproduct of autoxidation and peroxidation of fish and other oils (599).

II. Short Term Studies

Reports of fish oil toxicity in animals center on products of oxidation and deficiencies of vitamin E. These can end in damage to growth mechanisms, muscular dystrophy, multiple sclerosis, and brain damage. In these cases, however, the material fed would probably not have satisfied any regular quality standards for edible hydrogenated

fish oils, and the reports do not discuss extrapolation of the data to man. No reports of chronic fish oil toxicity in man have been found.

Dam (202) studied the production of exudative diathesis and of encephalomalacia as symptoms of vitamin E deficiency in chicks. He found that both symptoms were accelerated by feeding highly unsaturated long-chain FA; the source of such FA determined the relative prominence of the symptoms, and so did the protein/carbohydrate ratio of the diet. Addition of 1% of cholesterol accelerated the appearance of exudates when the diet contained 5% cod liver oil, and counteracted encephalomalacia when the diet contained 30% lard. Dam concluded that tissue damage, and not merely loss of antioxidant properties in the diet, caused the symptoms. He failed to reproduce either of the symptoms in rats, rabbits or ducklings; however, rabbits developed muscular dystrophy when fed a diet deficient in vitamin E.

Gershoff and Norkin (286) fed purified diets with various levels of vitamin E, with or without 5% tuna oil, to immature cats for up to 13.5 months. Cats fed tuna oil and no vitamin E developed steatitis with low levels of linoleic acid in their adipose tissue. Vitamin E supplements protected the cats completely against steatitis. When neither vitamin E nor tuna oil was added to the diet, mild symptoms of vitamin E deficiency, including muscle changes, developed after 12 months. These findings were interpreted in terms of the dietary requirements of the cat.

Harris and Embree (333) studied the relationship between intakes of PUFA and requirements for vitamin E in man. They calculated the average intakes of PUFA in grams per head for the population of the

United States for the year 1960; the average intakes of vitamin E were expressed as mg of d- α -tocopherol. The ratio, mg of vitamin E to g of PUFA, was found to be 0.6, and the authors inferred from published data that this ratio was probably marginal for protection of man against vitamin E deficiency. Fish oils were minimally involved in this study. Very recently the National Academy of Sciences has issued a denial of suggestions that vitamin E deficiency may be widespread in the United States at the present time (657).

Njaa et al. (668) included hydrogenated marine fat (HMF) (mainly from herring oil, unhydrogenated herring oil, or other fats) as 50% of the calories in a diet containing starch, sucrose, and albumin (10% w/w). These diets were fed to 29 groups of month-old rats for four weeks. Rats fed HMF grew relatively slowly and developed enlarged livers not due to fat, while rats fed the unhydrogenated herring oil grew better and developed fatty livers. The authors attributed the results of HMF to the presence of 20- and 22-carbon FA "of low saturation" together with trans and positional isomerization of the FA, in the HMF as fed.

Blaxter et al. (113) fed recently born bull calves a milk diet plus amounts of cod liver oil up to 4 oz per day for up to 71 days. Some calves died during the experiment; others were killed at the end. Autopsies revealed severe dystrophy of skeletal and heart muscles, but no lesions in livers, kidneys, or nervous systems. Partial protection was given by supplements of vitamins A and D.

Blaxter et al. (111) confirmed and amplified these findings "clinically, chemically, on dissection and histologically." They showed that the cause was the total unsaturated FA fraction of the

cod liver oil. No dystrophy was caused by the saturated FA fraction, the less unsaturated portion of the PUFA fraction, or the unsaponifiable residues of the cod liver oil. Only slight dystrophy was caused by the more unsaturated portion of the PUFA fraction. The tocopherol content of muscle samples did not differ as between normal and dystrophic muscles. No peroxidation of body fats was found. The authors concluded that the dystrophy was a general effect of the total PUFA fraction of the cod liver oil, and was unconnected with any hypervitaminosis A or D.

Subsequently Blaxter et al. (112) found that DL- α -tocopheryl acetate, given orally but not when given parenterally, protected similar calves against muscular dystrophy from cod liver oil supplements. Daily administration of 1 g of methylene blue gave similar protection and did not affect tissue levels of tocopherols. Ascorbic acid, ethyl gallate, or biotin gave no protection. The authors came to no conclusions about the mechanisms of these effects, but suggested a connection with oxidative phosphorylation.

A different picture emerged from four experiments on rats reported by Ershoff (251). Diarrhea and retarded growth resulted from supplementation of a low-fat, purified diet with crude or refined oils of tuna, sardine, menhaden, or cod liver. Substantial protection was given by additional supplements of cottonseed, soybean, sesame, corn, or wheat germ oils, by α -tocopherol, DPPD (N,N'-diphenyl-p-phenylenediamine), or Santoquin (6-ethoxy-2,2,4-trimethyl-1,2-dihydroquinoline), or by alfalfa meal, dessicated liver N.F., Torula yeast, casein, fish meal, DL-methionine, or a mixture of amino acids.

No protection was given by olive, coconut, or hydrogenated cottonseed oils; by butter fat or lard; by methyl linoleate; or by α -tocopheryl acetate or phosphate. The author drew no conclusions except in the introduction to his paper, where he remarked that "toxic effects may occur following fish oil administration due to factors other than the induction of a vitamin E-deficiency state."

Many investigators, in addition to Matsuo (see Acute Toxicity), have shown that toxic effects of oxidized oils can be attributed to peroxides and not to carbonyl derivatives.

Andrews et al. (58) studied the toxicity to rats of peroxides formed by autoxidation of soybean oil. They found that the peroxides were destroyed in the intestine, and concluded from this and other indications that the intestine was the site of toxic action. Inhibition of intestinal xanthine oxidase activity by autoxidized soybean oil and its reversal by exogenous FAD cofactor suggested interference with intestinal enzyme activities as a possible mechanism of peroxide toxicity. In these experiments this toxicity was demonstrated when the peroxide value of the added oil was as low as 100.

Rasheed et al. (739) fed rats a diet supplemented with 10% of pure triglycerides from fresh menhaden oil, and all was well; but when the triglycerides were oxidized experimentally to peroxide values of 125-310 they caused steatitis, enlarged livers, and high malonaldehyde (carbonyl) levels in blood and excreta. When the fresh oil was allowed to autoxidize for 24 hours and then was fed as 15% of the diet, it produced anorexia, steatitis, low hemoglobin,

and death, depending on the degree of autoxidation. Symptoms were largely reversed when rats, fed oil autoxidized to a peroxide value of 60, were treated with ethoxyquin or α -tocopherol; the rats were protected when these antioxidants were added to the fresh oil, but α -tocopheryl acetate, for example, was less effective than DL- α -tocopherol. High carbonyl levels in the blood were considered to be "a useful diagnostic criterion" for toxic effects of peroxidized oils.

However, Carpenter et al. (163) found no depression of growth in chicks fed de-fatted herring meal supplemented with herring oil pre-oxidized to a peroxide value of 142.

Roubal and Tappel (760) studied the effects of peroxidizing lipids, typified by ethyl arachidonate, upon the integrity of proteins in vitro. They found that transient free radicals, generated in the reaction mixture, damaged 17 different amino acids in proteins that included gamma-globulin, catalase, bovine serum albumin, ovalbumin, and hemoglobin. In addition, trypsin, pepsin, ovalbumin, α -chymotrypsin, cytochrome-c, and hemoglobin were rendered at least partly insoluble, and cysteine was partly converted to cystine with liberation of H_2S . This damage, according to the authors, "appears to be about one-tenth as effective as radiation damage."

Dam (204) illustrated the pathway by which peroxidic free radicals are generated during autoxidation of linoleic acid.

Reviewing these effects of autoxidation of fish oils, Stansby (832) emphasized that "These are all only potential effects. Whether or not they develop depends to a very large extent upon circumstances.... The most important factor influencing the degree to which these potential

alterations may develop is the degree and extent to which the poly-unsaturates oxidize. This in turn is largely controlled by the use to which the oil is put." These remarks were directed to all fish oils, not especially to hydrogenated fish oils, in which the degree of poly-unsaturation is, of course, diminished.

There appears to be a theoretical possibility that epoxides may arise either during metabolism of peroxides in the intestine or by action of drug-metabolizing enzyme systems on fish-oil hydrocarbons, and a further possibility that such epoxides may be carcinogenic. Grover (303) has reviewed the mechanisms. Apparently epoxides that form at the K regions (regions of high electron density associated with isolated phenanthrenoid double bonds) or other regions of cyclic hydrocarbons, are electrophilic. When these epoxides react with nucleic acids in vitro or in cultured cells, alkylation may occur; alternatively, an epoxide may be intercalated into DNA or may delete a DNA base, and either case will result in a frameshift mutation. Resultant morphologically abnormal cell colonies apparently have been found to be malignant on reinjection into rodents.

No reports have been found in which these effects have been studied in relation to hydrogenated fish oils in vitro or in vivo.

On the other hand, the rate of growth of carcinosarcoma in rats was reported by Haven (345) as slower when the dietary fat was cod liver oil of iodine number 9, than when it was coconut oil of iodine number 157. The difference was attributed to the difference in saturation of the FA.

III. Long Term Studies

No long term studies of chronic toxicity of hydrogenated fish oils have been found in the literature.

(Questions about hydrogenated PUFA, cholesterol deposition, and atherogenesis are outside the scope of this monograph. The present consumption of hydrogenated fish oils in the United States is virtually nil. Furthermore, the roles of saturated FA and blood cholesterol level in atherogenesis are matters of controversy, and cannot be debated here.)

IV. Special Studies

Pesticide residues. Pesticides of current concern that appear in sea fish comprise mainly the DDT family (DDT, DDE, and TDE). Over 5 ppm in fish or fractions of fish is unacceptable. Other substances that cause concern because of accidental contamination are the polychlorinated biphenyls (PCB). DDT and PCB are both toxic and persistent because they enter the food chain and stay there.

In 1968 Stout (848) monitored levels of the DDT family in several species of food fish in the Northwest Pacific and found levels up to 0.4 ppm; she attributed these accumulations to agricultural run-off by way of the Columbia River. In 1970 Stout et al. (849) reviewed these and other findings; for example, fish taken off Southern California had contained 20-50 ppm rising to 1,026 ppm, mainly in fat-rich tissues. Experiments had been undertaken to find ways of eliminating these residues; otherwise it would be useless to go fishing. It was found that all detectable amounts of p,p'-DDT in fish oils could be removed by refining, and 90% could be removed by hydrogenation; thus the regular processes (Figure 2) would normally eliminate all detectable

p,p'-DDT. However, p,p'-DDE, although substantially reduced, was not eliminated by these procedures. Stout et al. (849) concluded that the deodorization step was critical because temperatures were kept low (190-200°C) to avoid thermal degradation of the oil; therefore this step should be prolonged when pesticides of the DDT family were present or suspected.

These studies substantially confirmed earlier work by Smith et al. (816) on the removal of DDT, DDE, endrin, aldrin, dieldrin, heptachlor and heptachlor epoxide from vegetable oils.

Less work has been reported on PCB in fish oils. Beezhold and Stout (094) have described a gas chromatographic method of detecting PCB in fishery products. Problems have been encountered with standards, and studies are in progress. No reports have been found on the elimination of PCB from fish oils.

BIOCHEMICAL ASPECTS

I. Breakdown

Only oxidative changes seem to have been reported. Thermal degradation is not considered because hydrogenated fish oils have not been reported as used for deep-frying.

Possible toxic effects of oxidation have been discussed in foregoing sections. In addition, four sorts of flavor changes have been reported (828): (1) rancidity, (2) reversion to the "fishy" flavor of the original oil, (3) a "hardening" flavor resulting from hydrogenation and removed afterwards (see Figure 2), and (4) reversion to the hardening flavor.

These changes reportedly affect consumer acceptance, and much of the chemistry is obscure. The Codex Alimentarius recommendations (181) aim to define a product that is free from such defects. No reports have been found of tests using hydrogenated fish oils that specifically comply with those recommendations. Currently there is no significant use of hydrogenated fish oils in the food supply of the United States (see Consumer Exposure). The following data are mentioned because they may become relevant in the future.

Chipault (174) isolated volatiles from oxidized fish oils under vacuum at 100°C, by vacuum-steam distillation, and by adsorption from nitrogen gas onto charcoal. Paper chromatography of the acidic fractions revealed only acetic and propionic acids. The basic fractions contained small amounts of amines that were not identified. The carbonyl fractions contained formaldehyde, acetaldehyde,

propionaldehyde, butyraldehyde, valeraldehyde, hexaldehyde, acetone, butanone, pentanone, α - β -unsaturated monocarbonyls, glyoxal, other low-mw α -keto aldehydes and diketones, and 2,4-dinitrophenylhydrazones not further identified.

Wyatt and Day (948) isolated and measured quantitatively 26 carbonyl compounds from salmon oil autoxidizing under controlled conditions.

Meijboom (620) tried to relate the odors and tastes of some pure saturated and unsaturated aldehydes to their molecular structures. He found, as have others, that the technical problems were enormous; to date they remain unsolved (164). Nevertheless, he and his colleagues (458) appear to have identified the principal carrier of hardening flavor in hydrogenated linseed and soybean oils as 6-trans-nonenal, which was detected by some subjects at the concentration of only 0.3 ppb.

Hannewijk (324) has argued that if Keppler et al. are correct, 6-trans-nonenal should also carry the hardening flavor in hydrogenated fish oils. He has attributed the reversion of hardening flavor in hydrogenated fish oils to slight increases of bound aldehydes resulting from oxidation during storage.

According to Stansby (828), the stability and flavor of hydrogenated fish oils used in foods until 1951 presented problems; however, use of menhaden oil should cause fewer problems because it is less highly unsaturated. Menhaden oil is now, and foreseeably, the chief source of hydrogenated fish oils in the United States (see Consumer Exposure). Stansby adds that flavor reversion problems have been overcome with hydrogenated soybean oils, that fish oils contain similar ω -3 PUFA,

and that hydrogenated herring oils are used in the European food supply.

II. Absorption-Distribution

No indications were found in the literature that pathways of absorption and distribution of hydrogenated fish oils might differ from those of other dietary fats. In principle, ingested fats are digested to monoglycerides and FFA by bile action in the intestine; short-chain FA are absorbed into the bloodstream; and FA with chains longer than 8-12 carbons enter the lymph, diffuse into the thoracic duct, and filter gradually into the circulation. Hydrogenated fish oils are especially rich in long-chain FA (see Table 2).

Fats that are excreted in the feces normally have not been absorbed. Thus McCay and Paul (607) found that guinea pigs utilized about 60% of a hydrogenated vegetable oil, compared with 90% of unhydrogenated oils including salmon oil.

Fish oils polymerized by hydrogenation were reported by Matsuo (585,586,599) to damage the intestinal mucosa when fed to rats as sole sources of dietary lipids.

Beare (090) fed rats a variety of oils including partly hydrogenated herring oil and measured the FA distribution in liver, carcass, and milk fats. She found that the quantities of palmitic, palmitoleic, stearic, oleic, linoleic, linolenic, eicosenoic, eicosadienoic, and erucic acids in milk reflected the dietary pattern of FA, generally more than did the carcass FA. Tissue arachidonic acid, not provided in the diet, was related to dietary linoleic acid level. Changes of diet fat were followed by corresponding changes in tissue distribution

of long chain FA. Beare observed that the rats took a long time to adapt to utilization of erucic acid.

It is well known that some fatty acids, that are first synthesized in plankton, can persist through the food chain.

Mohrhauer and Holman (636) fed 108 weanling rats a fat-free diet plus purified linoleic and linolenic acids in several proportions and they determined the FA composition of lipids in heart, liver and adipose tissue by gas chromatography. Tissue levels of arachidonate were inversely related to amounts of dietary linolenate, showing that the linolenate inhibited conversion of linoleate to arachidonate. In another study, the same authors (637) fed 184 weanling rats ethyl linoleate, ethyl arachidonate, or ethyl linolenate, and measured the FA composition of the liver. It was found to correspond well with the diet.

Neudorffer and Lea (660) fed turkeys a basal diet plus 2.5% of beef fat or anchovy oil, or 5% of anchovy oil. The basal diet contained 2.3% of lipid richer in linoleate than either of the supplements. Analysis of skin fat revealed that FA had been transferred from the diet with high, though varying, efficiency. Peroxide values of the skin fat after incubation showed that the unhydrogenated anchovy oil had greatly reduced the stability of the skin fat.

Imaichi et al. (388) used gas-liquid chromatography to study the FA composition of 210 samples of depot fat removed at surgery from the abdominal or thoracic walls of human patients. Ten of these patients were known to be unusually high consumers of fish oils, and their

depot fat contained approximately double the levels of highly unsaturated FA found in the average of the other samples.

III. Metabolism and Excretion

The normal metabolism of FA present in fish oils has been studied; for example, "families" of FA tend to repress the metabolism of other FA "families". Thus, Holman and his colleagues (637) (372) found that increases of the linolenate/linoleate ratio in the diet progressively inhibited conversion of linoleate to arachidonate. Brenner and Peluffo (128) and Mohrhauer et al. (638) have extended Holman's findings. However, in these studies highly refined oils derived from fish oils were fed to animals as the sole dietary lipids, and the data probably do not illustrate what happens when fish oils contribute in small part to normal human diets. Nor did these studies involve partly hydrogenated fish oils of uncertain FA composition.

Hartwell (338) digested a number of different fats with pancreatic lipase in vitro. She found that coconut oil was the most rapidly digested; followed by palm kernel oil and castor oil; then butter fat; and lastly a slow group including beef and hog fats, a number of vegetable oils, cod liver oil, and hydrogenated whale and arachis oils.

The term "essential fatty acids" (EFA) originally meant FA that alleviated certain skin symptoms and growth failures in laboratory animals fed experimental diets. Later, EFA were identified as PUFA with the first double bond in the ω -6 position, e.g., linoleic acid (836). Then Privett et al. (728) (730) showed that PUFA of the linolenate series, having the first double bond in the ω -3 position, possessed

EFA activity when fed as 10% or more of experimental diets; they also found that fish oils were not inferior to corn oil as sources of EFA activity. However, these fish oils were not hydrogenated.

The EFA activity of partly hydrogenated fish oils has been studied by Aaes-Jørgensen since 1954 and also by Beare (1961). The findings have been summarized by Aaes-Jørgensen (1961) as follows:

"However, only very few experiments have been made so far with partially hydrogenated fish oils as dietary fat.

"In a study on the effect of partially hydrogenated marine oils as the sole dietary fat to rats, hydrogenated oil was used in an amount of 7% and 28% of an otherwise fat-free diet to weanling rats for 14 weeks, Aaes-Jørgensen (1954). The higher percentage of hydrogenated herring oil (28%) decreased the growth rate significantly as compared with the 7% level. The growth rate of the latter animals was significantly poorer than that of animals on 7% or 28% non-hydrogenated herring oil. Further, on the diet with 28% partially hydrogenated herring oil the animals ceased gaining weight after six weeks on the diet and began to lose weight after nine weeks. After 14 weeks on this diet the rats were moribund. The dermal symptoms indicating EFA deficiency were very pronounced in all the animals reared on diets with 7% or 28% partially hydrogenated herring oil. Hematuria developed in 50% of the animals fed on 7%, and in 100% of the animals fed on 28% hydrogenated herring oil. At the time when these experiments were carried out, the alkali isomerization procedure was the only available technique for determination of the unsaturated fatty acids. Thus, rather limited information only could be obtained concerning the fatty acids in the dietary fats. However, it was beyond doubt that the contents of polyenoic acid, normal as well as conjugated, were low.

"Beare (1961) fed diets with 2% partially hydrogenated herring oil to male, weanling Wistar rats for six weeks. Gas chromatographic analysis of the oil indicated a low content of linoleic acid, the presence of odd-numbered fatty acids, and of several series of C_{20} acids. The rats fed on the hydrogenated herring oil consumed more food and gained more weight than did the animals fed on diets with similar amounts of corn oil, rapeseed oil, or margarine (possessing some marine oil). However, when all weight gains were adjusted for food consumption by covariance analysis, the diet containing corn oil appeared to be the most efficiently utilized, and the diet containing margarine the least; rapeseed and hydrogenated herring oil were intermediate.

"The livers of the rats fed partially hydrogenated herring oil were significantly heavier than those of the other rats. Despite the small quantity of dietary linoleic acid available to these rats, the level of octadecadienoic acid in liver and carcass fat was similar to

that of rats fed on rapeseed oil or margarine. The author suggests a preferential retention of the sparsely available linoleic acid. In some livers there appeared to be an isomer of the common linoleic acid. The weight of the 14-15 days old litters of mothers fed on hydrogenated herring oil was significantly smaller than those of mothers supplied with corn oil or margarine. The rat milk fatty acids generally reflected more closely the dietary pattern of fatty acids than did those of the tissues.

"The nutritive effects of partially hydrogenated fish oils are only vaguely known and present a most difficult area to explore because of the complexity of the unsaturated fatty acid pattern of such oils. However, further studies in this area are needed and no doubt will be undertaken as the analytical techniques become more refined."

Aaes-Jørgensen and Højlmer (008) fed partly hydrogenated herring oil to rats as the sole dietary fat. Spermatogenic tissue was totally impaired after 5 weeks, contrasted with 15 weeks on partly hydrogenated arachis oil, and 26 weeks on fat-free diet. These findings were attributed to EFA deficiency.

Other reports attribute losses of nutritive value in fish oils to autoxidation. For example, feed intake, weight gain, and hemopoiesis diminished with increasing oxidation of menhaden oil fed to rats (739) and pigs (681). Stansby (836) has interpreted these data and others already noted (202) (113) (111) (112) (286) in terms of EFA deficiency.

As already noted (see Description), careful refinement should not destroy antioxidant properties of the original fish oils; however, more can be added (181). Dam (204) reviewed the activities of a range of antioxidants. Harris and Embree (333) estimated that 0.6 mg of D- α -tocopherol per g of polyenoic acids should supply the required amount of antioxidant activity. However, Olcott (676) has pointed out that the antioxidant activity of α -tocopherol is not commensurate with its total vitamin E activity.

The impairment of vitamin activities in fish oils by part-hydro-

genation does not seem to have been studied quantitatively. Dubin and Funk (235) reported that part-hydrogenation of cod liver oil in the presence of Ni at temperatures not over 115°C did not destroy vitamin A and D activities. However, van der Steur (913) found destruction at the higher temperatures of modern refining processes, though vitamin E survived better than A or D. He pointed to risks of over-supplementation of finished oils with synthetic vitamins A, D, or E.

Paluszak (693) hydrogenated a vegetable oil to mp 59°C and iodine number 5, eliminating linolenate, linoleate, and arachidonate. He fed this oil as a diet supplement to groups of young rats, with and without added cholesterol and biotin (1% and 2 ppm respectively in the diet). Losses of weight and of depot fat during 11 weeks of feeding were attributed to practical deficiencies of biotin and of EFA associated with the extreme hydrogenation.

Nevertheless authors mostly agree that partly hydrogenated fish oils, properly prepared, are nutritionally adequate sources of dietary fat (902) (913) (936).

As already noted, fecal excretion of significant amounts of dietary fats indicates failure of absorption (607). It is common knowledge that a major route of excretion of fats is through sweat. However, most fats are utilized by the body as energy sources.

IV. Effects on Enzymes and Other Biochemical Parameters

Bernsohn and Stephanides (100) have argued that multiple sclerosis can be caused by PUFA deficiency that results in demyelination of the nervous system. Kishimoto et al. (468) analyzed slices of the left

cerebral hemispheres of patients who died of multiple sclerosis and found subnormal axon densities and ganglioside concentrations in the plaques, and also a fatty acid distribution in the glycerophospholipids of the plaques that resembled grey matter rather than adjacent white matter. The white matter resembled that from control slices. It should be noted that hydrogenated fish oils do not figure in these reports.

The extensive literature on cholesterol-lowering effects of fish oils has been reviewed by Stansby (836). Although various explanations have been proposed for the "superior cholesterol depressant effect of fish oil fatty acids," the mechanisms remain "obscure." All of these explanations concern FA of fish oil before hydrogenation; the relevance to hydrogenated fish oils is not discussed. Stansby (836) also regards as speculative reports that glyceryl ethers and non-FA components of fish oils, such as squalene, have desirable nutritional properties.

Grollman (301) and Reichsman (746) reported that fish oils have properties that can be used to lower blood pressure in patients with chronic hypertension, but which do not alter blood pressure in normotensives. These reports did not involve hydrogenated oils.

No reports have been found of any use of hydrogenated fish oils to study the well known inhibitory effects of FA on the activities of some carbohydrate-metabolizing enzymes.

V. Drug Interaction

No reports have been found of studies of drug interaction. Neither have reports been found to suggest that hydrogenated fish oils differ

from other lipids in capacity to carry fat-soluble vitamins A, D, E and K.

VI. Consumer Exposure Information

Statistics for 1971 (Table 7) disclose heavy United States trade in fish oils but "negligible" use of hydrogenated fish oils in foods. Reasons for this have been discussed by Stansby and Knobl (839) and Stansby (828). During World War I, Californian sardines (pilchards) became important as a food source, and state law restricted their uses to food. By 1920 hydrogenated fish oils were used in margarine in Europe, but laws restricted the production and sale of all margarine through much of the United States. In 1925, 120,000 lbs of hydrogenated sardine oil were used in margarine in the United States; this figure rose to 40 million lbs in 1940, when less than half of these amounts of menhaden oil were so used. Additional unrecorded amounts of hydrogenated fish oils were used in shortenings. By 1951, however, overfishing had destroyed the California sardine stock and no fish oils were made into margarine or shortening in the United States from that year until 1967, when some herring oil was imported from Iceland for use in shortenings. This practice was quickly discontinued for fear that labeling requirements would force its disclosure.

The present FDA standard of identity for oleomargarine (167) does not permit the use of fish oils. When this standard was enacted in 1955 no petitions were received for the inclusion of such fish oils, and they were therefore omitted by default. No evidence for or against the use of fish oils has been considered by the FDA--either up to that

Table 7

Fish Oils - Consumer Exposure Tables

Production*	1971	1970	1949
World production, millions tons:			
Marine, vegetable, animal oils	41.373	39.236	
Vegetable oils	20.505	19.545	
Marine oils	1.261	1.292	
Fish oils including liver oils	1.095	1.052	
USA total of fats and oils, millions lbs:			
Food use	11,009	11,161	6,419
Non-food use	5,383	5,675	3,735
USA production of shortenings,** millions lbs:	3,479	3,599	1,494
Imports into USA, millions lbs***	1971	1969	1965
Cod oil except liver oil	5.974	3.764	4.765
Herring oil except liver oil	1.470	0.055	0.771
Other fish oils, NSPF, except liver oils	0.076	0.124	0.002
Sperm whale oil·**** crude	41.437		
refined	4.115		

* These data were abstracted from Agricultural Statistics, 1972, US Government Printing Office, Washington, D.C. (897)

** Content of fish oils called "negligible"

*** These data were abstracted from US Imports for Consumption and General Imports, US Department of Commerce, Bureau of the Census, publications FT 246-71, FT 246-69, and FT 246-65, US Government Printing Office, Washington, D.C. (896)

**** Imports now (1973) illegal.

time or since. However, the recent internationally recommended Codex Alimentarius specifications for margarine (182) permit the use of fish oils; the present U.S. standard of identity for oleo-margarine (067) is about to be reconsidered by the FDA (066). If no petition is made to include the use of fish oils, the present law will continue to require their omission from the standard of identity. On the other hand, if a petition is presented, evidence will be called for, and the Select Committee on GRAS substances review of hydrogenated fish oils will be potentially relevant evidence.

No such legal restrictions apply to the use of hydrogenated fish oils in shortenings in the United States. However, any plant in which a food component is prepared must meet sanitation requirements that have been adopted by the FDA; according to Stansby (828) "few, if any, fish oil plants in this country meet such requirements". This stricture does not apply, of course, to fish oils imported into the U.S. from countries where plants have met the FDA sanitation requirements, for example, the herring oil that was imported from Iceland in 1967 (828).

Practical inhibitions on resumption of the use of fish oils for food purposes in the United States include problems of price, public image, and technology concerned with elimination of undesirable trace components and flavors (828). The elimination problems appear to be soluble (828). Public images are notoriously malleable, and the 1973 increases in the price of soybean oil are relevant to any future demand for reintroduction of fish oils. Some data on world trade in fish oils 1962-1967 are given by Stansby and Knob1 (839).

- 1 A.D. Little, Inc. 1956
Commercial uses for menhaden oil
In, Com. Fish. Rev.: Final Report to Fish and
Wildlife Service, U.S. Dept. of the Interior,
Contract No. 14-49-008-2830, July 25 19(4a):3-4
- 2 Aaes-Jorgensen, E. 1954
The role of fat in the diet of rats. 6. Influence
on growth of various fats in ordinary refined
state and after hydrogenation or polymerization
Copenhagen: Store Nordiske Videnskabsboghandel
- 3 Aaes-Jorgensen, E. 1964
Essential fatty acids
Physiol. Rev. 44:1-51
- 4 Aaes-Jorgensen, E. 1964
Hydrogenated oils and polyunsaturated fatty-acid
isomers in nutrition
Proc. Nutr. Soc. 20:456-463
- 5 Aaes-Jorgensen, E. 1965
Biochemical aspects of hydrogenated oils and
isomeric unsaturated fatty acids in nutrition
Nutr. Dieta 7:130-134
- 6 Aaes-Jorgensen, E. 1966
Unsaturated fatty acid isomers in nutrition
Nutr. Rev. 24:1-5
- * 7 Aaes-Jorgensen, E. 1967
Fish oils as a source of essential fatty acids
In, Fish Oils, Stansby, M.E. (Ed.), AVI
Publishing Co., Westport, Conn. pp. 300-324
- * 8 Aaes-Jorgensen, E., and G. Holmer. 1969
Essential fatty acid-deficient rats. I. Growth
and testes development. III. Distribution of
lipid classes in rat testes after feeding partly
hydrogenated oils
Lipids 4:501-506, 515-521
- 9 Aaes-Jorgensen, E., and O.S. Privett. 1959
Biochemistry of lipids
Pergamon Press, Oxford
- 10 Aaes-Jorgensen, E., and R.T. Holman. 1958
Essential fatty acid activity. I. Content of
polyenoic acids in testes and heart as an
indicator of EPA status
J. Nutr. 65:633
- 11 Aaes-Jorgensen, E., O.S. Privett, and R.T. Holman.
1959
Essential fatty acid activities of hydrocarbons
and alcohols analogous to linoleate and
linolenate
J. Nutr. 67:443-421
- 12 Aaes-Jorgensen, E., and H. Dam. 1954
The role of fat in the diet of rats. 2. Influence
of dietary fats on growth
Brit. J. Nutr. 8(3):285-289
- 13 Aarhus Oliefab AS. 1970
Cocoa butter substitute
Brit. Pat. 1,208,629 issued in 1970
- 14 Abu-Nasr, A.M., W.M. Potts, and R.T. Holman. 1954
Highly unsaturated fatty acids. II.
Fractionation by urea inclusion compounds
J. Am. Oil Chem. Soc. 31:16-20
- 15 Ackman, R.G. . 1963
Structural correlation of unsaturated fatty acid
esters through graphical comparison of gas-
liquid chromatographic retention times on a
polyester substrate
J. Am. Oil Chem. Soc. 40:558
- 16 Ackman, R.G. 1964
Structural homogeneity in unsaturated fatty acids
of marine lipids. A review
J. Fish. Res. Board Can. 21:247-254
- 17 Ackman, R.G., and J.C. Sipos. 1964
Pilchard oil: An analysis for component fatty
acids with particular reference to the C24 chain
length
J. Fish. Res. Board Can. 21:841-843
- 18 Ackman, R.G., and J.C. Sipos. 1965
Isolation of the saturated fatty acids of some
marine lipids with particular reference to
normal odd-numbered fatty acids and branched-
chain fatty acids
Comp. Biochem. Physiol. 45:445-456
- 19 Ackman, R.G., and P.M. Jangaard. 1963
On the occurrence of 4,7,10,13-16-
docosapentaenoic acid in saury (*Cololabis saira*)
oil
J. Fish. Res. Board Can. 20:1551-1552
- 20 Ackman, R.G., and P.M. Jangaard. 1963
Identification of the major polyunsaturated C16
acids of marine oils by GLC separation factors
on normal and organosilicone polyesters
J. Am. Oil Chem. Soc. 40:744-747
- 21 Ackman, R.G., and R.D. Burgher. 1963
Component fatty acids of the milk of the grey
(Atlantic) seal
Can. J. Biochem. Physiol. 41:2501
- 22 Ackman, R.G., and R.D. Burgher. 1964
Cod liver oil: Component fatty acids as
determined by gas-liquid chromatography
J. Fish. Res. Board Can. 21:319-326
- 23 Ackman, R.G., and R.D. Burgher. 1964
Cod roe: Component fatty acids as determined by
gas chromatography
J. Fish. Res. Board Can. 21:469-475
- 24 Ackman, R.G., L.R. Galloway, P.M. Jangaard, and
M.L. Hughes. 1962
Conversion of the free fatty acids of cod oil to
methyl esters in situ
J. Fish. Res. Board Can. 19:605-614
- 25 Ackman, R.G., P.M. Jangaard, R.D. Burgher, M.L.
Hughes, and W.A. MacCallum. 1963
Observations on the oil and component fatty acids
of the oil from Newfoundland capelin
J. Fish. Res. Board Can. 20:591-596
- 26 Ackman, R.G., P.M. Jangaard, R.J. Hoyle, and H.
Brockerhoff. 1964
Origin of marine fatty acids. I. Analyses of
fatty acids produced by the diatom *Skeletonema
costatum*
J. Fish. Res. Board Can. 21:747-756
- 27 Ackman, R.G., R.D. Burgher, and J.L. Hughes. 1962
The free fatty acids of cod oil. 4. Anomalous
composition by fatty acid chain length
J. Fish. Res. Board Can. 19:1061-1084
- * 28 Ackman, R.G., S.W. Hooper and J. Hingley. 1971
Monoethylenic fatty acids of a partially
hydrogenated herring oil
J. Am. Oil Chem. Soc. 48:804-806
- 29 Adachi, A. 1960
Cuttlefish XII. Existence of C24 acids, C26 acids
and C28 acids
J. Jap. Oil Chem. Soc. 9:591-593
- 30 Adamson, L.F. 1960
Serum cholesterol concentrations of various
ethnic groups in Hawaii
J. Nutr. 71:27-36
- 31 Adamstone, F.B. 1941
Cholesterol content of brain in nutritional
encephalomalacia of vitamin E deficient chicks
Arch. Pathol. 34:711
- 32 Agawa, T., Y. Hirao, and S. Komori. 1953
The liver oil of a cod, *Lotella phycis*
J. Jap. Oil Chem. Soc. 2:246-249
- 33 Agduhr, E. 1926
Post-natal development under different conditions
of nutrition and circumstances of functioning,
changes in heart through presence of cod liver
oil in food
Acta Paediatr. 5:319

- Agduhr, E. 1926
anges in the organism caused by cod liver oil
added to the food
Acta Paediatr. 6:165
- 35 Agduhr, E. 1928
Are the so called "A" vitamins in cod liver oil
the cause of its toxic effect on the organism;
and can a basal diet, complete as regards the so
called "B" and "C" vitamin content, prevent this
toxic effect?
Acta Paediatr. 7:289
- 36 Agduhr, E. 1936
Toxicity, influence on some organic structures of
different components
Z. Vitaminforsch. 5:27-37
- 37 Agduhr, E., G. Blix, and B. Vahlquist, 1934-1935
Studies on the influence of some natural fats and
their components on animal tissue structures.
II. The influence on some organic structures of
different components of cod liver oil
Upsala Lakareforen. 40(3/4):183-388
- 38 Ahrens, E.H. Jr., and L.C. Craig. 1952
Separation of the higher fatty acids
J. Biol. Chem. 195:299-310
- 39 Ahrens, E.H., J. Hirsch, W. Insull, and M.L.
Paterson. 1958
Chemistry of lipids as related to atherosclerosis
C.C. Thomas, Springfield
- 40 Ahrens, E.H., Jr., J. Hirsch, W. Insull, Jr.,
T.T. Tsaltus, R. Bloomstrand, and M.L. Peterson.
1957
The influence of dietary fats on serum lipid
levels in man
Lancet 1:943-953

Ahrens, E.H., Jr., W. Insull, Jr., J. Hirsch, W.
Stoffel, M.L. Peterson, J.M. Parquahar, T.
Miller, and R.J. Thomasson. 1959
Effect on human serum lipids of a dietary fat,
highly unsaturated, but poor in unsaturated
fatty acids
Lancet 1:115-119
- 42 Ajinomoto Co. Inc. 1962
Process of manufacturing fatty oils having low
melting point and no fishy odor from marine
animal fatty oil
Jap. Pat. 4021 issued Jun. 11, 1962
- 43 Albrink, M.J., and E.B. Mann. 1959
Serum triglycerides in coronary artery disease
Arch. Intern. Med. 103:4-8
- 44 Alexander, J.C. 1966
Effect on diet handling on nutritional studies
with used frying fats
Lipids 1:254-257
- 45 Alfin-Slater, R.B. 1961
Factors affecting essential fatty acids
utilization
In, Drugs Affecting Lipid Metabolism, Garattini
and Paoletti (Eds.), Elsevier Publ., Amsterdam
- 46 Alfin-Slater, R.B., L. Aftergood, A.P. Wells, and
H.J. Deuel. 1954
The effect of essential fatty acid deficiency on
the distribution of endogenous cholesterol in
the plasma and liver of the rat
Arch. Biochem. Biophys. 52:180-185
- 47 Allen, R.R. 1960
Hydrogenation
J. Am. Oil Chem. Soc. 37:521-523
- 48 Allen, R.R. 1962
Practical aspects of hydrogenation
J. Am. Oil Chem. Soc. 39:457-459
- * 49 Altman, P.L., and D.S. Dittmer (Eds.). 1972
Biology Data Book
Federation of American Societies for Experimental
Biology, Bethesda, Md. 1:300-352
- 50 American Medical Association, Council on Food and
Nutrition. 1962
The regulation of dietary fat
J. Am. Med. Assoc. 181:411
- 51 American Oil Chemists Society. 1959
Official and Tentative Methods of the American
Oil Chemists Society, 2nd Edition
American Oil Chemists Society, Chicago, Illinois
- * 52 American Oil Chemists Society. 1965
Official and Tentative Methods of the American
Oil Chemists Society, 10th Edition
American Oil Chemists Society, Chicago, Illinois
- * 53 American Pharmaceutical Association. 1970
National Formulary
Am. Pharm. Assoc., Washington, D.C. 13th Ed.
- 54 American Society for Testing Material. 1965
Book of ASTM Standards
Am. Soc. Testing Materials, Philadelphia, Pa.
Part 20:287
- 55 Analytical Methods Committee, Report prepared by
the Vitamin E Panel. 1959
The determination of tocopherols in oils, foods
and feeding stuffs
Analyst 84:356-372
- 56 Anderson, J.T., A. Keys, and F. Grande. 1957
The effects of different food fats on serum
cholesterol concentrations in man
J. Nutr. 62:421-444
- 57 Anderson, K.L., and M.A. Steinberg. 1964
Effect of lipid content on protein-sodium
linolenate interaction in fish muscle homogenates
J. Food Sci. 29:327-330
- * 58 Andrews, J.S., W.H. Griffith, J.P. Mead, and R.A.
Stein. 1960
Toxicity of air-oxidized soybean oil
J. Nutr. 70:199-210
- 59 Anglemier, A.F. 1958
Menhaden and pilchard oils as caloric supplements
for swine
Univ. Microfilms Pub., Diss. Abstr. 18,1190
No. 24622
- 60 Anglemier, A.F., and J.E. Oldfield. 1957
Feeding of various levels of pilchard oil to swine
J. Anim. Sci. 16:922-926
- 61 Anon. 1922
Fish by-products on the Pacific Coast
Pac. Fisherman 20(5):40 pp.
- 62 Anon. 1934, 1937, 1939
Fishy flavor in foodstuffs
Univ. of Reading, Natl. Inst. Res. Dairying Ann.
Repts. for years ending Sep. 30, 1934, Sep. 30,
1937, Sep. 30, 1939
- 63 Anon. 1949
Norsk fiskeri- og Fangsthandbok I. (Handbook of
Norwegian Fisheries I)
Cannermyer, Oslo
- 64 Anon. 1966
Firms manufacturing marine animal scrap, meal,
oil, solubles, and homogenized condensed fish,
1965
Statistical List, Curr. Fish. Statistics, U.S.
Dept. Int., Washington, D.C. No. 151 (revised)
- 65 Anon. 1968
Industrial Fishery Products-1967 Review:
Situation and Outlook, Current Economic Analysis
II, April 1968
Bureau of Commercial Fisheries, Fish and Wildlife
Service, U.S. Dept. of Interior, Washington
- * 66 Anon. 1973
Oleomargarine, margarine
Fed. Regist. 38(49):6968

- * 67 Anon. 1973
Code of Federal Regulations, Food and Drugs,
Oleomargarine, Margarine
Fed. Regist.
Title 21, Chap. I, Part 45, Amended to March 14
- 68 Arimune, K. 1951
The hydroxylation of oils and fats. I. Selective
hydroxylation of unsaturated fatty acids. II.
Partial hydroxylation of methyl linoleate
J. Chem. Soc. Jap., Ind. Chem. Sec.
54:739-741; 54:760-761
- 69 Asmundson, V.S., T.H. Jukes, H.M. Fyler, and M.L.
Maxwell. 1938
Effect on certain fish meals and fish oils in the
ration on the flavor of the turkey
Poult. Sci. 17:147-151
- * 70 Association of Official Agricultural Chemists.
1960
Official Methods of Analyses of the Association
of Official Agricultural Chemists
AOAC, Washington, D.C.
- 71 Association of Official Agricultural Chemists.
1965
Official methods of analysis
Association of Official Chemists, Washington, D.C.
10th Ed.
- 72 Association of Official Agricultural Chemists.
1965
International union of pure and applied chemistry
carbamate method.
Official Methods of Analysis of the Association
of Agricultural Chemists, AOAC, Washington, D.C.
24:023-24, 028
- 73 Aure, L. 1962
Fatty acid composition of food fats
J. Agric. Food Chem. 10:120-123
- 4 Aure, L. 1967
Manufacture of fish-liver oil
In, Fish Oils, M.E. Stansby (Ed.), AVI
Publishing, Westport, Conn. pp. 193-205
- 75 Ayres, E.B. 1958
Polyolefinic alcohols of increased conjugation
from soybean and menhaden oil
U.S. Pat. 2,819,317 issued Jan. 7, 1958
- 76 Bailey, A.E. 1945
Industrial oil and fat products
Interscience Press, New York City
- 77 Bailey, A.E. 1951
Industrial oil and fat products
Interscience Publishers, N.Y.
- 78 Bailey, B.E. 1952
The chemistry and technology of marine oils with
particular reference to those of Canada
Fish. Res. Board Can. Bull., Ottawa p. 89
- 79 Baker, R.W.R., R.H.S. Thompson, and K.J. Zilkha.
1963
Fatty-Acid composition of brain lecithins in
multiple sclerosis
Lancet 1:26-27
- 80 Banks, A. 1950
Some factors affecting the oxidation of the fat
of dehydrated herrings
J. Sci. Food Agric. 1:28-34.
- 81 Banks, A. 1952
The development of rancidity in cold-stored
herrings. The influence of some antioxidants
J. Sci. Food Agric. 3:250-256
- 82 Banks, A. 1967
Deteriorative changes in fish oils
In, Fish Oils, Stansby, M.E. (Ed.), AVI
Publishing Co., Westport, Conn. pp. 148-163
- 83 Barnes, R.H., W.O. Lundberg, H.T. Hanson, and
G.O. Burr. 1943
The effect of certain dietary ingredients on the
keeping quality of body fat
J. Biol. Chem. 149:313-322
- 84 Barry, H.C., J.C. Hundley, and L.Y. Johnson. 1965
Pesticide Analytical Manual, 2.21(8)
U.S. Department of Health, Education, and
Welfare, Food and Drug Administration
- 85 Barry, H.C., J.C. Hundley, and L.Y. Johnson. 1965
Pesticide Analytical Manual, 2.32
U.S. Department of Health, Education, and
Welfare, Food and Drug Administration
- 86 Bateman, L., H. Hughes, and A.J. Morris. 1953
Hydroperoxide decomposition in relation to the
initiation of radical chain reactions
Disc. Faraday Soc. 14:190-199
- 87 Bazalar, A. 1965
Technological study of the hydrogenization
process of fish oil
Faculty of Chemistry, Univ. of San Marcos Thesis
- * 88 Beal, R.E.; K.J. Moulton, H.A. Moser, and L.T.
Bleck. 1969
Removal of copper from hydrogenated soybean oil
J. Am. Oil Chem. Soc. 46(9):498-500
- 89 Beard, F.J., J.A. Schulz, and C.G. Culbertson.
1935
Effect of ingested menhaden and coconut oils upon
firmness of beef fat
Am. Soc. Anim. Pro. Rec. Proc. 28th Annu. Meet.
1935:286-290
- * 90 Beare, J.L. 1961
The influence of dietary fat on the fatty acid
composition of liver, carcass, and milk of rats
Can. J. Biochem. Physiol. 39:1855-1863
- 91 Beare, J.L. 1962
Fatty acid composition of food fats
J. Agric. Food Chem. 10:120-123
- 92 Becker, E.N. 1920
Zur verwendung gehärteter fette in der
speisefettindustrie
Chem. Ind. 2:63
- 93 Bedford, F., and C.E. Williams. 1911
Treating fish oil with hydrogen in the presence
of a catalyzer for the purpose of obtaining an
industrial, valuable fat, operating as specified
in 63,931
Swiss. Pat. 63,932 issued Dec. 18, 1911
- * 94 Beezhold, F.L., and V.F. Stout. 1973 (in press)
The use and effect of mixed standards on the
quantitation of polychlorinated biphenyls
Bull. Environ. Contam. Toxicol. 10(1)
- 95 Bell, E.R., J.H. Raley, F.F. Rust, F.H. Seubold,
and W.E. Vaughan. 1951
Reactions of free radicals associated with low
temperature oxidation of paraffins
Disc. Faraday Soc. 10:242
- 96 Bergmann, W., and R.A. Landowne. 1958
Contributions to the study of marine products.
XLVI. Phospholipids of a sea anemone
J. Org. Chem. 23:1241-1245
- 97 Bergmann, W., S.M. Creighton, and W.M. Stokes.
1956
Waxes and triglycerides of sea anemones
J. Org. Chem. 21:721-728
- 98 Bergstrom, S. 1945
On the oxidation of linoleic acid with lipoxidase
Ark. Kemi, Miner., Geol. 21A(15)
- 99 Bergstrom, S., H. Danielsson, and B. Samuelsson.
1964
The enzymatic formation of prostaglandin E2 from
arachidonic acid. Prostaglandins and related
factors 32
Biochem. Biophys. Acta 90:207-210

- * 100 Bernsohn, J., and L.M. Stephanides. 1967
Aetiology of multiple sclerosis
Nature 215:821-823
- 01 Bertram, S.H. 1937
Der nachweis tierischer fette und ole,
insbesondere geharteter tranen in fettgemischen
Ole, Fette, Wachse, Seife, Kosmet. 1937(2):13-14
- 102 Beveridge, J.M.R., W.F. Connell, and G.A. Mayer.
1956
Dietary factors affecting the level of plasma
cholesterol in humans: The role of fat
Can. J. Biochem. Physiol. 34:441-455
- 103 Beveridge, J.M.R., W.F. Connell, and G.A. Mayer.
1957
The nature of the substance in dietary fats
affecting the level of plasma cholesterol in
humans
Can. J. Biochem. 35:257-270
- 104 Beveridge, J.M.R., W.F. Connell, G.A. Mayer, and
H.L. Haust. 1958
Plant sterols, degree of unsaturation, and
hypocholesterolemic action of certain fats
Can. J. Biochem. Physiol. 36:895-911
- 105 Bickford, W.G., P. Krauczunas, and D.H. Wheeler.
1942
The reaction of nonconjugated unsaturated fatty
acid esters with maleic anhydride
Oil Soap 19:23-27
- 106 Biely, J., B.E. March, and D.A. Silvertrini. 1954
Fat studies in poultry. I. Herring Oil,
Santoworse-80, and Thyroprotein in the
laying ration
Poult. Sci. 33:1130-1135
- 107 Bittenbender, C.D. 1950
Gallic acid ester antioxidants for fish oils
U.S. Fish and Wildlife Service, Com. Fish. Rev.
12(2):1-18
- 108 Black, M.M., and H.M. Schwartz. 1950
South African fish products. XIX Seasonal
changes in the composition of pilchard (*Sardina
ocellata* Jenyns). XXXI. The composition of
pilchard oil and of masbanker oil
J. Sci. Food Agric. pp. 183-185, 248-251
- 109 Blank, M.L., L.J. Nutter, and O.S. Privett. 1966
Determination of the structure of lecithins
Lipid 1:132-135
- 110 Blaxter, K.L. 1962
Muscular dystrophy in farm animals: its cause
and prevention
Proc. Nutr. Soc. 21:211-216
- * 111 Blaxter, K.L., F. Brown, and A.M. MacDonald. 1953
The nutrition of the young Ayrshire calf 13. The
toxicity of the unsaturated acids of cod-liver
oil.
Brit. J. Nutr. 7:287-298
- * 112 Blaxter, K.L., F. Brown, W.A. Wood, and A.M.
MacDonald. 1953
The nutrition of the young Ayrshire calf 14.
Some effects of natural and synthetic anti-
oxidants on the incidence of muscular dystrophy
induced by cod-liver oil
Brit. J. Nutr. 7:337-349
- * 113 Blaxter, K.L., W.A. Wood, and A.M. MacDonald.
1953
The nutrition of the young Ayrshire calf 11. The
toxicity of cod-liver oil
Brit. J. Nutr. 7:34-50
- 114 Bligh, E.G., and W.J. Dyer. 1959
A rapid method of total lipid extraction and
purification
Fish. Res. Board Can., Studies 552-586:911-917
- 115 Bligh, E.G., and W.S. Dyer. 1959
A rapid method of total lipid extraction and
purification
Can. J. Biochem. Physiol. 37:911-917
- 116 Bloch, J. 1964
The importance of unsaturation to life
In, Metabolism and Physiological Significance of
Lipids. R.M.C. Dawson and D.M. Rhodes (Eds.),
John Wiley and Sons, London
- 117 Block, K., and D. Rittenberg. 1942
The biological formation of cholesterol from
acetic acid
J. Biol. Chem. 143:297-298
- 118 Blomstrand, R., and E.J. Ahrens, Jr. 1959
Absorption of chylol alcohol in man
Proc. Soc. Exp. Biol. Med. 100:802
- 119 Bodman, J., and J.H. Maisin. 1958
The alpha-glyceryl ethers
Clin. Chim. Acta 3:253-274
- 120 Boekennoogen, H.A. 1964
Analysis and characterization of oils, fats, and
fat products
Interscience Publishers, London
- 121 Boelhouwer, C., J. Gerckens, O.T. Lie, and H.I.
Waterman. 1953
Displacement of the double bonds during
hydrogenation of unsaturated fatty acid methyl
esters
J. Am. Oil Chem. Soc. 30:59-61
- 122 Bottino, N.R., and R.R. Brenner. 1962
Evidence for fat secretion in the intestine of
the fish
J. Am. Oil Chem. Soc. 39:519-520
- 123 Bozian, R.C., and J.G. Coniglio. 1962
The effect of ethyl arachidonate upon the
metabolism of oleic acid in the fat deficient rat
Fed. Proc. 21:286
- 124 Bozian, R.C., and J.G. Coniglio. 1963
The effect of arachidonate upon eicosatrienoate
levels in livers of fat-deficient rats
Fed. Proc. 22:489
- 125 Braekkan, O.R., G. Lambertsen, P. Utne, and L.R.
Njaa. 1968
Hydrogenated marine fat, its influence on the
fatty acid composition of depot fats and liver
lipids in the rat
Nutr. Diet. 10:24-44
- 126 Braamsnaes, F. 1964
Fiskemel og fiskeolje. (Fish meal and fish oil)
"Danmarks Fiskerierhverv" I, Liber A/S
- 127 Braun, G. 1929
Oxidation of unsaturated compounds. I. Oxidation
of crotonic and isocrotonic acids, of the
lactone of 3-hydroxyisocrotonic acid and of
maleic anhydride
J. Am. Chem. Soc. 51:228-248
- * 128 Brenner, R.R., and R.O. Peluffo. 1966
Effect of saturated and unsaturated fatty acids
on the desaturation in vitro of palmitic,
stearic, oleic, linoleic, and linolenic acids
J. Biol. Chem. 241:5213-5219
- 129 Brenner, R.R., D.V. Vazza, and M.E. De Tomas.
1963
Effect of a fat-free diet and of different
dietary fatty acids (palmitate, oleate, and
linoleate) on the fatty acid composition of fresh-
water fish lipids
J. Lipid Res. 4:341-345
- 130 British Standards Institution. 1958
Methods of analysis of oils and fats
British Standard 684:1958, British Standard
House, 2, Park St., London W.1Y. 4AA.

- 131 Brockerhoff, H. 1965
A stereospecific analysis of triglycerides
J. Lipid Res. 6:10-15

Brockerhoff, H., and R.J. Hoyle. 1963
On the structure of the depot fats of marine fish
and mammals
Arch. Biochem. Biophys. 402:452-455
- 133 Brockerhoff, H., and R.J. Hoyle. 1965
Hydrolysis of triglycerides by the pancreatic
lipase of a skate
Biochem. Biophys. Acta 98:435-436
- 134 Brockerhoff, H., M. Yurkowski, R.J. Hoyle, and
R.G. Ackman. 1964
Fatty acid distribution of lipids of marine
plankton
J. Fish. Res. Board Can. 21:1379-1384
- 135 Brockerhoff, H., R.G. Ackman, and R.J. Hoyle.
1963
Specific distribution of fatty acids in marine
lipids
Arch. Biochem. Biophys. 100:9-12
- 136 Brockerhoff, H., R.J. Hoyle, and K. Ronald. 1964
Retention of the fatty acid distribution pattern
of a dietary triglyceride in animals
J. Biol. Chem. 239:735-739
- 137 Brocklesby, H.N., and K. Green. 1934
Method for production of fish liver oils from
livers of low content
Fish. Res. Board Can. Rep. 22:18
- 138 Brohult, A. 1958
Effects of alkoxy glycerols and especially
selachyl alcohol on the bone marrow
in connection with irradiation treatment and in
leukemia therapy
Nature 181:1484-1485

Brohult, A. 1963
Alkoxyglycerols and their use in radiation
treatment
Acta Radiol., Suppl. 223:7-96
- 140 Bronte-Stewart, B., A. Keys, and J.F. Brock. 1955
Serum cholesterol, diet, and coronary heart
disease. An interracial survey in the Cape
Peninsula
Lancet 2:1103-1107
- 141 Bronte-Stewart, B.A., A. Antonis, L. Eales, and
J.F. Brock. 1956
Effects of feeding different fats on serum
cholesterol levels
Lancet 1:521-530
- 142 Brown, J.B. 1931
The nature of the highly unsaturated fatty acids
stored in the lard from pigs fed on menhaden oil
J. Biol. Chem. 90:133-139
- 143 Brown, J.B., and D.K. Kolb. 1955
Applications of low temperature crystallization
in the separation of the fatty acids and their
compounds
In, Progress in the Chemistry of Fats, R.T.
Holman, W.O. Lundberg, and T. Malkin (Eds.)
Pergamon Press, New York
- 144 Brown, J.B., and T.S. Sutton. 1931
Effect of feeding menhaden fish oil on secretion
of milk and composition of butterfat in the
dairy cow
J. Dairy Sci. 14:125-135
- 145 Brown, W.D., and A.L. Tappel. 1959
Fatty acid oxidation by carp liver mitochondria
Arch. Biochem. Biophys. 85:449-158
- 146 Brown, W.D., A.W. Venolia, A.L. Tappel, H.S.
Olcott, and M.E. Stansby. 1957
Oxidative deterioration in fish and fishery
products. II. Progress on studies concerning
mechanism of oxidation of oil in fish tissues
Commer. Fish. Rev. 19(5A):27-31
- 147 Bunnell, R.H., L.D. Matterson, E.P. Singsen, and
H.D. Eaton. 1956
Studies on encephalomalacia in the chick. V. The
effect of fish oil and diphenyl-p-
phenylenediamine on the vitamin metabolism of the
chick
Poult. Sci. 35:436-451
- 148 Bunnell, R.H., L.D. Matterson, E.P. Singsen, L.M.
Potter, A. Kozeff, and E.L. Jungherr. 1954
Studies on encephalomalacia in the chick. III.
The influence of feeding or injecting various
tocopherols or other antioxidants on the
incidence of encephalomalacia
Poult. Sci. 33:1046
- 149 Bunnell, R.H., L.D. Matterson, E.P. Singsen, L.M.
Potter, A. Kozeff, and E.L. Jungherr. 1955
Studies on encephalomalacia in chicks. 3. The
influence of feeding or injecting various
tocopherols and other antioxidants on the
incidence of encephalomalacia
Poult. Sci. 34:1068-1075
- 150 Burdick, G.E. 1964
The accumulation of DDT in lake trout and the
effect on reproduction
Trans. Am. Fish. Soc. 93(2):127-136
- 151 Burgess, J.F., and J.E. Pritchard. 1948
Tocopherol (vitamin E) therapy in sclerosis of
the legs with ulcer
Can. Med. Assoc. J. 59:242-247
- 152 Burr, G.O., and M.M. Burr. 1929
A new deficiency disease produced by the rigid
exclusion of fat from the diet
J. Biol. Chem. 82:345-367
- 153 Burr, G.O., and M.M. Burr. 1930
On the nature and role of the fatty acids
essential in nutrition
J. Biol. Chem. 86:587-621
- 154 Burr, G.O., M.M. Burr, and W.R. Brown. 1930-1931
On the nutritive value of certain oils
Proc. Soc. Exp. Biol. Med. 28:905-907
- 155 Burton, D.G., and L.F. Byrne. 1954
The constitution of sulphated oils, Part IV-
Effects of variations in the method of
manufacture on the composition of a sulphated
fish oil
J. Soc. Leather Trades' Chem. 38:10-22
- 156 Butler, C. 1949
The fish liver oil industry
U.S. Fish and Wildlife Service Fishery Leaflet
No. 233
- 157 Butterfield, R.O., and H.J. Dutton. 1964
Monitoring counter-current distribution with a
recording refractometer
Anal. Chem. 36:903-906
- 158 Butterfield, R.O., C.R. Scholfield, and H.J.
Dutton. 1964
Preparation of 9,15-octadecadienoate isomers
J. Am. Oil Chem. Soc. 41:397-400
- 159 Buxton, L.O. 1942
Effect of carbon treatment of fish liver oils,
vitamin A destruction and peroxide formation
Ind. Eng. Chem. 34:1486
- 160 Calcagna, O. 1935
Treatment of leprosy with oils obtained from some
salt- and fresh-water fishes
Sem. Med. B. Aires 11:557-562
- 161 Callow, E.H., and C.H. Lea. 1939
The effect of olive oil and cod liver oil in the
pig's diet on the quality of pork, sausages,
bacon and hams, and the influence of diet on the
susceptibility of the pig's fat to oxidation
Rep. Food Invest. Board (Gr. Brit.) for 1939
pp. 35-38

- 162 Carlson, W.E. 1966
Some nutritional studies of the naturally occurring alpha glyceryl ethers
Rep. Pac. Agric. Univ. Brit. Columbia, Vancouver
B.C. 128 pp.
- 163 Carpenter, K.J., C.H. Lea, and L.J. Parr. 1963
Chemical and nutritional changes in stored herring meal. 4. Nutritional significance of oxidation of the oil
Brit. J. Nutr. 17:151-169
- 164 Carr, C.J., H.F. Sassoon, and K.D. Fisher. 1973
A study of the potential applications of olfactory research in man
LSRO, PASEB for Department of the Army, Washington, D.C. Contract DAHC19-72-C-0010
- 165 Carrick, C.W., and S.M. Hauge. 1926
The effect of cod liver oil upon flavor in poultry meat
Poult. Sci. 5:213-215
- 166 Carroll, K.K. 1961
Quantitative estimation of peak areas in gas-liquid chromatography
Nature 191:377-378
- 167 Century, B., and M.K. Horwitt. 1959
Effect of fatty acids on encephalomalacia
Proc. Soc. Exp. Biol. Med. 102:375-377
- 168 Century, B., and M.K. Horwitt. 1969
Role of diet lipids in the appearance of dystrophy and creatinuria in the vitamin E deficient rat
J. Nutr. 72:357
- 169 Century, B., L.A. Witting, C.C. Harvey, and M.K. Horwitt. 1961
Composition of skeletal muscle lipids of rats, fed diets containing various oils
J. Nutr. 75:341-346
- 170 Century, B., M.K. Horwitt, and P. Bailey. 1959
Lipid factors in the production of encephalomalacia in the chick
Arch. Gen. Psychiatr. 1:420
- 171 Chang, H.-Y., and S. Gladstone. 1965
Separation of oleic acid and linoleic acid by solvent extraction
J. Am. Oil Chem. Soc. 42:346-347
- 172 Chang, S.S. 1967
Processing of fish oils
In, Fish Oils, Stansby, M.E. (Ed.), AVI Publishing Co., Westport, Conn. pp. 206-221
- 173 Chipault, J. 1960
The odors of oxidized fish oils
Annu. Rep., Hormel Inst., 1959-1960 pp. 7-8
- 174 Chipault, J.R. 1959
The odors of oxidized fish oils
Ann. Rept. Hormel Inst., 1958-1959 pp. 7-8
- 175 Chipault, J.R. 1962
High energy irradiation
In, Symposium on Foods: Lipids and Their Oxidation, H.W. Schultz (Ed.), AVI Publishing Co., Westport, Conn.
- 176 Chipault, J.R., and E. McMeans. 1965
Volatile acids from menhaden oil
J. Agric. Food Chem. 13:15-17
- 177 Christensen, P.K., and N.A. Sorensen. 1951
Studies related to pristane. V. The constitution of zameene
Acta Chem. Scand. 5:751-756
- 178 Clingman, A.L., D.E.A. Rivett, and D.A. Sutton. 1954
The chemistry of polymerized oils. Part IV. Thermal polymerization of some long-chain polyunsaturated fatty esters
J. Chem. Soc. 1954:1088-1090
- 179 Codex Alimentarius Commission. 1969
General standard for the labelling of pre-packaged foods
Codex Alimentarius Commission, Food and Agriculture Organization of the United Nations Sections 1,2,4, and 6. Ref. CAC/RSI-1969
- 180 Codex Alimentarius Commission. 1969
Determination of soap content
Methods of Analysis for Edible Fats and Oils, Codex Alimentarius Commission, Food and Agriculture Organization of the United Nations CAC/RM 13-1969
- * 181 Codex Alimentarius Commission. 1969
Recommended International General Standard for Edible Fats and Oils
Codex Alimentarius Commission, Food and Agriculture Organization of the United Nations, World Health Organization CAC/RS 19-1969
- * 182 Codex Alimentarius Commission. 1969
Recommended International Standard for Margarine
Codex Alimentarius Commission, Food and Agriculture Organization of the United Nations, World Health Organization CAC/RS 32-1969
- 183 Codex Alimentarius Commission. 1969
Determination of iron content
Methods of Analysis for Edible Fats and Oils, Codex Alimentarius Commission, Food and Agriculture Organization of the United Nations CAC/RM 14-1969
- 184 Coenen, J.W.E. 1960
Rate studies of the selective hydrogenation of fatty oils
Actes Congr. Int. Catal., Paris II:2705-2731
- 185 Coenen, J.W.E., H. Boerema, B.G. Linsen, and B. De Vries. 1964
Selectivity in fatty oil hydrogenation. Influence of catalyst pore structure on selectivity characteristics
Proc. 3rd Int. Congr. Catal., Amsterdam
- 186 Coffin, D.L., and J. Holzworth. 1954
"Yellow fat" in two laboratory cats: acid-fast pigmentation associated with a fish-base ration
Cornell Vet. 44:63
- 187 Coleman, J.E., C. Ricciuti, and D. Swern. 1956
Improved preparation of 9,10; 10,9-ketohydroxystearic acids by oxidation of oleic acid with potassium permanganate in neutral solution
J. Am. Chem. Soc. 78:5342-5345
- * 188 Committee on Specifications. 1972
Food Chemicals Codex
Committee on Food Protection, National Academy of Sciences, National Research Council, Washington, D.C. 2nd Edition
- 189 Common, R.H., E.W. Crampton, F. Farmer, and A.S.W. De Prietas. 1957
Studies to determine the nature of the damage to the nutritive value of menhaden oil from heat treatment
J. Nutr. 62:341-347
- 190 Cooke, S.R.B. 1958
Progress in the studies of utilization of fish oil derivatives in ore flotation
Comm. Fish. Rev. 20(1):14-19
- 191 Corbin, E.A., D.P. Schwartz, and M. Keeney. 1960
Liquid-liquid partition chromatography separation of the 2,4-dinitrophenylhydrazones of saturated aldehydes, methyl ketones, 2-enals and 2,4-dienals
J. Chromatog. 3:322
- 192 Cordy, D.R. 1954
Experimental production of steatitis (yellow fat disease) in kittens fed a commercial, canned cat food and prevention of the condition by vitamin E
Cornell Vet. 44:310

193 Cordy, D.R., and C.J. Stillinger. 1953
Steatitis ("yellow fat disease") in kittens
North Am. Vet. 34:714

Cormier, M. 1948
Antivitamin action of fish liver oils
Bull. Soc. Chim. Biol. 30:921-940

195 Cousins, E.R., W.A. Guice, and R.O. Feuge. 1959
Positional isomers formed during the
hydrogenation of methyl linoleate under
various conditions
J. Am. Oil Chem. Soc. 36:24-28

196 Cowan, J.C. 1962
Dimer acids
J. Am. Oil Chem. Soc. 39:534-545

197 Crampton, E.W., F.A. Farmer, and F.M. Berryhill.
1951
The effect of heat treatment on the nutritional
value of some vegetable oils
J. Nutr. 43:431-440

198 Crossley, A., T.D. Heyes, and B.J.P. Hudson. 1962
The effect of heat on pure triglycerides
J. Am. Oil Chem. Soc. 39:9

199 Cubero, J.M., and H.K. Mangold. 1965
Chromatography on adsorbent layers impregnated
with silver nitrate
Microchem. J. 9:227-236

200 Dalgaard-Mikkelsen, S., H.C. Momberg-Jorgensen,
and P.H. Petersen. 1958
Control of yellow fat disease in mink
Beretn. Forsgslab. Kobenhavn. 308:31 pp.

201 Dam, H. 1943
Effect of cod liver oil and rancidity on certain
vitamin E deficiency symptoms
Proc. Soc. Exp. Biol. Med. 52:285-287

Dam, H. 1944
Studies on vitamin E deficiency in chicks
Nutr. 27:193-211

203 Dam, H. 1949
Relationship of vitamin E deficiency to tissue
peroxides
Ann. N.Y. Acad. Sci. 52:195

* 204 Dam, H. 1957
Influence of antioxidants and redox substances on
signs of vitamin E deficiency
Pharmacol. Rev. 9:1-16

205 Dam, H., and H. Granados. 1945
Peroxidation of body fat in vitamin E deficiency
Acta Physiol. Scand. 10:162-171

206 Dam, H., and J. Glavind. 1938
Alimentary exudative diathesis
Nature, London 142:1077

207 Dam, H., and J. Glavind. 1939
Alimentary exudative diathesis, a consequence of
E-avitaminosis
Nature 143:810

208 Dam, H., and J. Glavind. 1939
Alimentary exudative diathesis and its relation
to vitamin E
Skand. Arch. Physiol. 82:299

209 Dam, H., and J. Glavind. 1942
Factors influencing capillary permeability in the
vitamin E deficient chick
Science 96:235

210 Dam, H., G.K. Nielsen, I. Prange, and E.
Sondergaard. 1958
Exudative diathesis produced by vitamin E
deficient diets without polyenoic fatty acids
Experientia 14:291-292

211 Dam, H., H. Granados, and I. Prange. 1949
Influence of two different fats on reproduction
capacity of vitamin E deficient rats
Acta Physiol. Scand. 18:161

212 Dam, H., I. Kruse, I. Prange, and E. Sondergaard.
1951
Substances affording a partial protection against
certain vitamin E deficiency symptoms
Acta Physiol. Scand. 22:299-310

213 Dam, H., J. Glavind, O. Bernth, and E. Hagens.
1938
Anti-encephalomalacia activity of dl-alpha-
tocopherol
Nature, London 142:1157

214 Dam, R., C. Kristensen, C.K. Nielsen, and E.
Sondergaard. 1959
Influence of dietary cholesterol, cod liver oil
and linseed oil on cholesterol and polyenoic
fatty acids in tissues from fasted and non-fasted
chicks
Acta Physiol. Scand. 45:31-42

215 Dansky, L.M. 1962
The growth promoting properties of menhaden fish
oil as influenced by various fats
Poult. Sci. 41:1352-1354

216 Davies, A.W., and T. Moore. 1941
Interaction of vitamins A and E
Nature, London 147:794-796

217 Davies, W.L. 1939
"Fishiness" as a flavor and taint
Flavours 2:18-21

218 Davies, W.L., and E. Gill. 1936
Investigations on "fishy" flavors
Chem. Ind. (London) 55:141T-146T

219 Davis, C.L., and J.R. Gorham. 1954
The pathology of experimental and natural cases
of "yellow fat" disease in swine
Am. J. Vet. Res. 15:55

220 Davis, G.K., and L.A. Maynard. 1938
Cod liver tolerance in calves
J. Dairy Sci. 21:145-152

221 De Groot, A.P., and S.A. Reed. 1959
Influence on dietary cod-liver oil and some
fractions of cod-liver oil on serum cholesterol
levels of rats
Nature 183:1191

* 222 De Konig, A.J. and M.H. Silk. 1963
The 2-thiobarbituric acid reagent for
determination of oxidative rancidity in fish oils
J. Am. Oil Chem. Soc. 40(5):165-169

223 De Konig, A.J. 1963
The phospholipids of pilchard and hake
Fish. Ind. Res. Inst., Ann. Rep. 17:64

224 Deblois, E.T. 1882
The origin of the menhaden industry
Bull. of the U.S. Fish Comm., Vol. 1 for 1881
1:46-51

* 225 Deck, R.E., and K.K. Kaiser. 1970
Analytical method for determining copper in
edible shortenings and oils
J. Am. Oil Chem. Soc. 47:126-128

226 Deuel, H.J., Jr. 1951
The Lipids, Their Chemistry and Biochemistry
Interscience Publishers, New York Vol. I

227 DeSesa, R.J. 1959
Fish oil in modern coatings
Paint, Oil Chem. Rev. 122(5):6-10

228 DeVries, B. 1963
Quantitative separations of higher fatty acid
methyl esters by adsorption chromatography on
silica impregnated with silver nitrate
J. Am. Oil Chem. Soc. 40:184-186

- 229 DeWitt, K.W. 1963
Seasonal variations in cod-liver oil
J. Sci. Food Agric. 14:92-98
- 230 DiTullio, N.W., C.S. Jacobs, Jr., and W.L. Holmes. 1965
Thin-layer chromatography and identification of free sterols
J. Chromatog. 20:354
- 231 Dolev, A. and H.S. Olcott. 1965
The triglycerides of sable fish (*Anaplocoma fimbria*). I. Quantitative fractionation by column chromatography on silica gel impregnated with silver nitrate
J. Am. Oil Chem. Soc. 42:624-627
- 232 Dolev, A., and H.S. Olcott. 1965
The triglycerides of sable fish (*Anaplocoma fimbria*). II. Fatty acid distribution in triglyceride fractions as determined with pancreatic lipase
J. Am. Oil Chem. Soc. 42:1046-1051
- 233 Domart, C., D.T. Miyauchi, and W.N. Sumerwell. 1955
The fractionation of marine-oil fatty acids with urea
J. Am. Oil Chem. Soc. 32:481-483
- 234 Dreosti, G.M. 1967
Fish Oil Industry in South Africa and Southwest Africa
In, Fish Oils, M.E. Stansby (Ed.), AVI, Westport, Conn. pp. 428-432
- * 235 Dubin, H.E. and C. Funk. 1923
Studies on the chemistry of cod liver oil. I. The effect of hydrogenation upon the vitamin content
Proc. Soc. Exp. Biol. Med. 21:139-141
- 236 Dunning, E.T. 1959
Drying oils in printing inks
J. Am. Oil Chem. Soc. 36:586-590
- 237 Dutton, H.J. 1963
Kinetics of linolenate hydrogenation
J. Am. Oil Chem. Soc. 40:35
- 238 Dutton, H.J., C.R. Scholfield, E.P. Jones, E.H. Pryde, and J.C. Cowan. 1963
Hydrazine-reduced linolenic acids as a source of C9, C12, and C15 dibasic acids
J. Am. Oil Chem. Soc. 40:175-179
- 239 Dyer, J.A. 1967
General industrial and potential uses of fish oils
In, Fish Oils, Their Chemistry, Technology, Stability, Nutritional Properties, and Uses, M.E. Stansby (Ed.), The AVI Publishing Company, Inc., Westport, Conn. 19:270-279
- 240 Eckey, E.W. 1948
Process for treating fats and fatty oils
U.S. Pat. 2,442,531 issued Jun. 1, 1948
- 241 Eckey, E.W. 1948
Treatment of glycerides for use in edible fats
U.S. Pat. 2,442,532 issued Jun. 1, 1948
- 242 Eckey, E.W. 1948
Treatment of glycerides
U.S. Pat. 2,442,533 issued Jun. 1, 1948
- 243 Eckey, E.W. 1949
Directed interesterification in glycerides
Ind. Eng. Chem. 40:1183-1190
- 244 Edwards, H.M., Jr., and J.E. Marion. 1963
Influence of dietary menhaden oil on growth rate and tissue fatty acids of the chick
J. Nutr. 81:123-130
- 45 Edwards, H.M., Jr., J.E. Marion, and J.C. Driggers 1962
Fat and fatty acid requirements of poultry
Proc. 12th World's Poult. Congr. pp. 182-186
- 246 Einset, E., H.S. Olcott, and M.E. Stansby. 1957
Oxidative deterioration in fish and fish products. IV. Progress on studies concerning oxidation of extracted oils
U.S. Fish and Wildlife Service, Com. Fish. Rev. 19(5a):35-37
- 247 Ellis, C. 1930
Hydrogenation of organic substances
3rd ed., pp. 22, 184, 866
- 248 Emker, E.A., C.R. Scholfield, and H.J. Dutton. 1964
Chromatographic separation of cis and trans fatty esters by argentation with a macroreticular exchange resin
J. Am. Oil Chem. Soc. 41:388-390
- 249 Emmerie, A., C. Engel, and W. Klip. 1952
The tuberculostatic action in vitro of the unsaponifiable fraction of cod liver oil
J. Sci. Food Agric. 3:264-268
- 250 Erikson, A., R.H. Coots, F.H. Mattson, and A.M. Kligman. 1963
Effect of P/S ratio, partial hydrogenation of dietary fats, and dietary cholesterol upon plasma cholesterol in man
Circulation 28:656
- * 251 Ershoff, B.H. 1960
Effects of diet on fish oil toxicity in the rat
J. Nutr. 71:45-53
- 252 Esso Research and Engineering Company. 1959
Lubricating grease and process of their manufacture
Ger. Pat. 1,054,631 issued Mar. 12, 1959
- 253 Etinburg, Je., and L. Nikolejewa. 1936
Hydrierung von tran
Oel- Fett-Ind. 12:212-214
- 254 Evans, H.M., and G.A. Emerson. 1943
The prophylactic requirement of the rat for alpha tocopherol
J. Nutr. 26:555
- 255 Evans, R.S., and P.G.W. Scott. 1961
Precision and accuracy in gas chromatographic analysis
Nature 190:710-712
- 256 Farkas, T., and S. Herodek. 1962
Origin of the characteristic fatty acid composition of aquatic organisms
Magy. Tud. Akad. Tihanyi Biol. Kutatoint, Evkonyve 29:79-83
- 257 Farkas, T., and S. Herodek. 1964
The effect of environmental temperature on the fatty acid composition of crustacean plankton
J. Lipid Res. 5:369-373
- 258 Farmer, E.H. 1946
Peroxidation in relation to olefinic structure
Trans. Faraday Soc. 42:228-236
- 259 Farmer, F.A., B.C. Mutch, J.M. Bell, L.D. Woolsey, and E.W. Crampton. 1950
The vitamin E requirement of guinea pigs
J. Nutr. 42:309
- 260 Farquhar, J.W., W. Insull, P. Rosen, W. Stoffel, and E.H. Ahrens. 1959
The analysis of fatty acid mixtures by gas-liquid chromatography
Nutr. Rev. Aug.(Suppl.):17
- 261 Feuge, R.O., E.R. Cousins, S.P. Fore, E.F. DuPre, and R.T. O'Connor. 1953
Modification of vegetable oils. XV. Formation of isomers during hydrogenation of methyl linoleate
J. Am. Oil Chem. Soc. 30:454-460
- 262 Filer, L.J., Jr., R.E. Rumery, and K.E. Mason. 1946
Specific unsaturated fatty acids in the production of acid-fast pigment in the vitamin E-deficient rat and the protective action of tocopherols
In, Biological Antioxidants, Trans. of the First Conf., Oct. 10-11, C.G. Mackenzie, (Ed.), N.Y., Josiah Macy, Jr. Foundation p. 67

- 263 Fineberg, H., and A.G. Johanson. 1967
Industrial use of fish oils
In, *Fish Oils, Their Chemistry, Technology, Stability, Nutritional Properties, and Uses*, M.E. Stansby (Ed.), The AVI Publishing Company, Inc., Westport, Conn. (Also, reprinted in 1967 as U.S. Fish and Wildlife Service, Circular 278, no separate pagination-) pp. 222-238
- 264 Fitzpatrick, J.D., and M. Orchin. 1958
Fate of selenium in the isomerization of oleic acid
J. Org. Chem. 23:918-919
- 265 Folsch, J., I. Ascoli, J.A. Meath, and F.N. LeBaron. 1951
Preparation of lipid extracts from brain tissue
J. Biol. Chem. 191:833-841
- 266 Fontell, J., R.T. Holman, and G. Lambertsen. 1960
Some new methods for separation and analysis of fatty acids and other lipids
J. Lipid Res. 1:391-404
- * 267 Food and Agriculture Organization. 1973
FAO Books in Print
Food and Agriculture Organization of the United Nations pp. 66-69
- 268 Fore, S.P., and W.G. Bickford. 1959
Catalytic hydrogenation of 9,10-epoxyoctadecanol and 9,10-epoxyoctadecyl acetate
J. Org. Chem. 24:620-622
- 269 Fore, S.P., and W.G. Bickford. 1961
Catalytic hydrogenation of cis-6,7-epoxyoctadecanoic acid
J. Org. Chem. 26:2104-2105
- 270 Forss, D.A., E.A. Dunstone, and W. Stark. 1960
Fishy flavor in dairy products. III. The volatile compounds associated with fishy flavor in washed cream
J. Dairy Res. 27:373
- 271 Forss, D.A., E.A. Dunstone, and W. Stark. 1960
The volatile compounds associated with tallowy and painty flavors in butterfat
J. Dairy Res. 27:381
- 272 Forss, D.H., E.A. Dunstone, and W. Stark. 1960
Fishy flavor in dairy products. II. The volatile compounds associated with fishy flavor in butter fat
J. Dairy Res. 27:211-220
- 273 Frankel, E.N., C.D. Evans, and J.C. Cowan. 1960
Thermal dimerization of fatty ester hydroperoxides
J. Am. Oil Chem. Soc. 37:418-424
- 274 Fraser, E.B., J.H. Stothart, and H.S. Gutteridge. 1934
Fish meals and oils: Feeding value for livestock and poultry
Dom. Can. Dept. Agric. Pamphlet 163:7-10
- 275 Freeman, S.F. 1940
Process of extracting and refining glycerides and products resulting therefrom
U.S. Pat. 2,200,390 issued May 14, 1940
- 276 Fulco, A.J., and J.P. Mead. 1959
Metabolism of essential fatty acids. VIII. Origin of 5,8,11-eicosatrienoic acid in the fat-deficient rat
J. Biol. Chem. 234:1411
- 277 Fulco, A.J., and J.P. Mead. 1960
The biosynthesis of the octadecadienoic acids in the rat
J. Biol. Chem. 235:3379-3384
- 278 Gaddis, A.M., Rex Ellis, and G.T. Currie. 1960
Carbonyls in oxidizing fat. III. The distribution of volatile and nonvolatile carbonyls
Food Res. 25:495
- 279 Gaddis, A.M., Rex Ellis, and G.T. Currie. 1961
Carbonyls in oxidizing fat. V. The composition of neutral volatile monocarbonyl compounds from autoxidized oleate, linoleate, linolenate esters and fats
J. Am. Oil Chem. Soc. 38:371
- 280 Gaddis, A.M., Rex Ellis, and G.T. Currie. 1961
Girard T reagent for carbonyls
Nature 191:1391
- 281 Gambal, D., and F.W. Quackenbush. 1960
Effects of cholesterol and other substances on essential fatty acid deficiencies
J. Nutr. 70:497-501
- 282 Gast, L.E., E.W. Bell, and H.M. Teeter. 1956
Reactions of conjugated fatty acids. III. Kinetics of the Diels-Alder reaction
J. Am. Oil Chem. Soc. 33:278-281
- 283 Gauglitz, E.J., Jr., and D.C. Malins. 1960
The preparation of polyunsaturated aliphatic aldehydes via the acyloin condensation
J. Am. Oil Chem. Soc. 37:425-427
- 284 Gauglitz, E.J., Jr., and E.H. Gruger, Jr. 1965
Adsorptive bleaching and molecular distillation of menhaden oil
J. Am. Oil Chem. Soc. 42:561-563
- 285 Gellerman, J.L., and H. Schlenk. 1959
Column chromatography of lipids - odd number straight chain fatty acids of menhaden oil
Experientia 15:387-388
- * 286 Gershoff, S.N., and S.A. Morkin. 1962
Vitamin E deficiency in cats
J. Nutr. 77:303-308
- 287 Glavind, J., H. Granados, S. Hartmann, and H. Dam. 1949
A histochemical method for the demonstration of fat peroxides
Experientia 5:84-85
- 288 Glavind, J., S. Hartmann, J. Clemmesen, K.E. Jessen, and H. Dam. 1952
Studies on the role of lipoperoxides in human pathology. II. The presence of peroxidized lipids in the atherosclerotic aorta
Acta Path. Microbiol. Scand. 30:1-6
- 289 Gloor, U., F. Weber, and O. Wiss. 1961
Vitamin A and the biosynthesis of cholesterol
In, *Drugs Affecting Lipid Metabolism*, Garattini and Paoletti (Eds.), Elsevier Publ., Amsterdam
- 290 Goettsch, M. 1942
Alpha-tocopherol requirement of the mouse
J. Nutr. 23:543
- 291 Golumbic, C., and H.A. Mattill. 1940
The oxidation of vitamin E
J. Biol. Chem. 134:535-541
- 292 Gordon, H.H., H.M. Nitowski, and M. Cornblath. 1955
Studies of tocopherol deficiency in infants and children
Am. J. Dis. Child. 90:669
- 293 Gorham, J.R., G.A. Baker, and N. Boe. 1951
Observations on the etiology of yellow fat disease in mink
Vet. Med. 46:100
- 294 Gorham, J.R., N. Boe, and G.A. Baker. 1951
Experimental "yellow fat" disease in pigs
Cornell Vet. 41:332
- 295 Graham, W.R., Jr., and P.T. Cupps. 1938
Action of herring oil before and after hydrogenation on yield and fat percentage of milk of the goat
J. Dairy Sci. 21:45-48

- 296 Granados, H., and H. Dam. 1950
On the histochemical relationship between peroxidation and the yellow-brown pigment in the adipose tissue of vitamin E deficient rats
Acta Path. Microbiol. Scand. 27:591-596
- 97 Green, D.E., and S.J. Wakil. 1960
Enzymatic mechanisms of fatty acid oxidation and synthesis
In, *Lipid Metabolism*, K. Bloch (Ed.), John Wiley and Sons, N.Y. and London
- 298 Greenberg, D.M., C.E. Calbert, E.E. Savage, and H.J. Deuel. 1950
The effect of fat level of the diet on general nutrition. VI. The interrelation of linoleate and linolenate in supplying the essential fatty acid requirement in the rat
J. Nutr. 41:473
- 299 Greenspan, P.P., and R.J. Gall. 1956
Epoxy fatty acid ester plasticizers. Preparation and properties
J. Am. Oil Chem. Soc. 33:391-394
- 300 Griffiths, T.W. 1961
Studies on the requirement of the young chick for vitamin E. II. The effects of different sources and levels of dietary lipids live-weight gain and body vitamin E storage
Brit. J. Nutr. 15:271
- * 301 Grollman, A. 1945
Preparation of extracts from oxidized marine and other oils for reducing blood pressure in experimental and human chronic hypertension
J. Pharmacol. Exp. Ther. 84:128-135
- 302 Groot, E.H., and H.J. Klein Obbink. 1953
Influence of heating cod-liver oil to 230 °C on growth of young rats
Voeding 14:123-126
- 303 Grover, P. 1973
How polycyclic hydrocarbons cause cancer
New Scientist 58(850):685-687
- 304 Gruger, E.H. 1960
Fractionation and purification of triglycerides, fatty acids, and methyl esters from fish oils
Proc. Gulf-Caribbean Fish. Inst. 13th Session pp. 53-57
- * 305 Gruger, E.H., Jr. 1967
Fatty acid composition
In, *Fish Oils*, Stansby, M.E. (Ed.), AVI Publishing Co., Westport, Conn. pp. 3-30
- 306 Gruger, E.H., Jr., D.C. Malins, and E.J. Gauglitz, Jr. 1960
Glycerolysis of marine oils and the preparation of acetylated monoglycerides
J. Am. Oil Chem. Soc. 37:214-217
- 307 Gruger, E.H., Jr., R.W. Nelson, and M.E. Stansby. 1964
Fatty acid composition of oils from 21 species of marine fish, freshwater fish, and shellfish
J. Am. Oil Chem. Soc. 41:662-667
- 308 Gumbmann, M., W.D. Brown, and A.L. Tappel. 1958
Intermediate metabolism of fishes and other aquatic animals
U.S. Fish and Wildlife Service Special Scientific Report--Fisheries 288:28-31
- 309 Gunstone, F.D. 1958
An Introduction to the Chemistry of Fats and Fatty Acids
John Wiley and Sons, New York
- 310 Gutkin, S.S. 1950
Miscellaneous uses of drying oils
J. Am. Oil Chem. Soc. 27:538-544
- 311 Guyer, K.E., W.A. Hoffman, L.A. Horrocks, and D.G. Cornwell. 1963
Studies on the composition of glyceryl ethers and their preparation from diacyl glyceryl ethers in liver oils
J. Lipid Res. 4:385-391
- 312 Hadorn, H., and K. Zurcher. 1970
Universal-methode zur gas-chromatographischen untersuchung von speisefetten und oelen. (Trennung der fettsaure-methylester von C4 bis C24 und quantitative bestimmung des milchfettes in beliebigen fettsauremischungen). (Universal method for gas chromatographic examination of edible fats and oils. (Separation of C4-C24 fatty acid methyl Dtsch. Lebensm.-Rundsch. 66(3):77-87
- 313 Haighton, A.J. 1959
The measurement of the hardness of margarine and fats with cone penetrometers
J. Am. Oil Chem. Soc. 36:345-348
- 314 Haighton, A.J. 1963
Consistency of margarine and fats
Fette, Seifen, Anstrichm. 65:479-482
- 315 Haighton, A.J. 1965
Work-softening of margarine and shortening
J. Am. Oil Chem. Soc. 42:27-30
- 316 Haissinsky, M. (Ed). 1958
Les peroxydes organiques en radiobiologie
Masson, Paris
- 317 Hakansson, I. 1966
Physico-chemical changes in artificial fat emulsions during storage
Acta Chem. Scand. 20:2267-2281
- 318 Hallgren, B., and S. Larsson. 1962
The glyceryl ethers in the liver oils of elasmobranch fish
J. Lipid Res. 3:31-38
- 319 Hallgren, B., and S.O. Larsson. 1959
Separation and identification of alkoxyglycerols
Acta Chem. Scand. 13:2147-2148
- 320 Hammond, E.G., and W.O. Lundberg. 1953
A methyl docosahexaenoate: Its isolation and characterization
J. Am. Oil Chem. Soc. 30:438-441
- 321 Hammond, E.G., and W.O. Lundberg. 1953
The alkali isomerization of a methyl docosahexaenoate and the spectral properties of conjugated fatty acids
J. Am. Oil Chem. Soc. 30:433-438
- 322 Hammond, J.C. 1941
A factor in cod liver oil that hinders the utilization of vitamin E by chicks
Poult. Sci. 20:369-371
- 323 Hanahan, D.J., H. Brockeroff, and E.J. Barron. 1960
The site of attack of phospholipase (lecithinase) A on lecithin: A re-evaluation
J. Biol. Chem. 235:1917-1993
- * 324 Haanewijk, J. 1967
Use of fish oils in margarine and shortening
In, *Fish Oils*, Stansby, M.E. (Ed.), AVI Publishing Co., Westport, Conn. pp. 251-269
- 325 Hansen, K., H.T. Offerdahl, and B. Eriksen. 1933
Resorption of hardened whale fat in human intestine
Norsk Mag. Laegevidensk. 94:1305-1317
- 326 Hansley, V.L. 1935
Preparation of high molecular weight acylolins
J. Am. Chem. Soc. 57:2303-2305
- 327 Hansley, V.L. 1947
Sodium reduction of fatty acid esters
Ind. Eng. Chem. 39:55-62
- 328 Hanson, N.W. 1964
Chemical and nutritional aspects of oxidized and heated fats
Chem. Ind. 1964:1541-1549

- 329 Hanson, S.W.F., and J. Olley. 1963
Application of Bligh and Dyer method of lipid extraction to tissue homogenates
Biochem. J. 89(3):101P
- 330 Hardin, J.O., J.L. Milligan, and V.D. Sidwell. 1964
The influence of solvent extracted fish meal and stabilized fish oil in broiler rations on performance and on the flavor of broiler meat
Poult. Sci. 43:858-860
- 331 Harlow, C.M., and A.R. Morton. 1962
Use of fish in the control of hypercholesterolemia and obesity
In, Fish in Nutr., Heen and Kreuzer (Eds.), Fishing News Books, London pp. 292-293
- 332 Harris, L.A., and H.S. Olcott. 1966
Reaction of aliphatic tertiary amines with hydroperoxides
J. Am. Oil Chem. Soc. 43:11-14
- * 333 Harris, P.L., and W.D. Embree. 1963
Quantitative consideration of the effect of polyunsaturated fatty acid content of the diet upon the requirements for vitamin E
Am. J. Clin. Nutr. 13:385-392
- 334 Harris, P.L., E.G. Hardenbrook, P.P. Dean, E.R. Cusack, and J.L. Jensen. 1961
Blood tocopherol values in normal human adults and incidence of vitamin E deficiency
Proc. Soc. Exp. Biol. Med. 107:381
- 335 Harris, P.L., M.L. Quaife, and W.J. Swanson. 1950
Vitamin E content of foods
J. Nutr. 40:367
- 336 Hart, P.A. 1958
Fish oil research at the Seattle Fishery Technological Laboratory
Com. Fish. Rev. 20(4):17-21
- 337 Hartmann, S., and J. Glavind. 1949
A new sensitive method for the determination of peroxides of fats and fatty acids
Acta Chem. Scand. 3:954-958
- * 338 Hartwell, G.A. 1938
IX. A note on the digestion of fats by pancreatic lipase
Biochem. J. 32:462-466
- 339 Harwood, H.J. 1954
Nitrogen-containing derivatives of the fatty acids
J. Am. Oil Chem. Soc. 31:559-563
- 340 Harwood, H.J. 1962
Reactions of the hydrocarbon chain of fatty acids
Chem. Rev. 62:99-154
- 341 Hashimoto, T., K. Nukada, H. Shiina, and T. Tsuchiya. 1963
On the structure of highly unsaturated fatty acids of fish oils by high resolution nuclear magnetic resonance spectral analysis
J. Am. Oil Chem. Soc. 40:124-128
- 342 Hass, G.M. 1938
Intercellular transformations of unsaturated fatty acids and esters
Arch. Path. 26:1196
- 343 Hatano, M., K. Zama, and H. Igarashi. 1961
Lipids of salmonoid fishes. I. Acetone-soluble lipid from liver of salmon, *Oncorhynchus masu*
Bull. Jap. Soc. Sci. Fish. 27:1001-1004
- 344 Hauge, J.G., and R. Nicolaysen. 1959
The serum cholesterol depressive effect of linoleate, linolenic acids and of cod liver oil in experimental hypercholesterolemic rats
Acta Physiol. Scand. 45:26-30
- * 345 Haven, F.L. 1936
The effect of cod-liver oil on tumor growth
Am. J. Cancer 27:95-98
- 346 Heard, D.H., T. Moore, and I. Sharman. 1952
The dialuric acid haemolysis test for vitamin E deficiency in rats
Biochem. J. 52:1
- 347 Heaton, F.W., and N. Uri. 1958
Improved iodometric methods for the determination of lipid peroxides
J. Sci. Food Agric. 9:781-786
- 348 Heesch, A. 1963
Modern margarine making
Fette, Seifen, Anstrichm. 65:125-137, 205-214
- 349 Hegsted, D.M., A. Gotsis, and F.J. Stare. 1957
The effects of various fats upon experimental hypercholesterolemia in the rat
J. Nutr. 63:377-391
- 350 Heller, J.H., M.S. Heller, S. Springer, and E. Clark. 1957
Squalene content of various shark livers
Nature 179:919-920
- 351 Henderson, C., W.L. Johnson, and A. Inglis. 1969
Organochlorine insecticide residues in fish (National Pesticide Monitoring Program)
U.S. Congr. Record-Senate (June 9, 1969), Pestic. Monit. J. pp. 56142-56157; 3(3):145-171
- 352 Henick, A.S., M.F. Benica, and J.H. Mitchell. 1954
Estimating carbonyl compounds in rancid fats and foods
J. Am. Oil Chem. Soc. 31:88
- 353 Henly, R.S. 1965
Preparative gas-liquid chromatography of lipids
J. Am. Oil Chem. Soc. 42:673-681
- 354 Herb, S.P., L.P. Witnauer, and R.W. Riemenschneider. 1951
Isolation of eicosapentaenoic and docosapentaenoic acids from natural sources as their methyl esters by adsorption and distillation techniques
J. Am. Oil Chem. Soc. 28:505-507
- 355 Herb, S.P., R.W. Riemenschneider, and J. Donaldson. 1951
Isolation of natural arachidonic acid as its methyl ester
J. Am. Oil Chem. Soc. 28:55-58
- 356 Herstad, O. 1970
Effect of fat-supplements in broiler feed
Meld. Norg. Landbr. Hoysk. 49:1-55
- 357 Heyden, R., G. Dieckelmann, and J. Plapper. 1960
Oiling medium for the manufacture of whiter and lighter colored leather and pelts for fur
Ger. Pat. 973,930 issued Jul. 28, 1960
- 358 Hilditch, T.P., and P.N. Williams. 1964
The Chemical Constitution of Natural Fats
John Wiley and Sons, New York 4th Ed.
- 359 Hilditch, T.P., and P.N. Williams. 1964
The chemical constitution of natural fats
Chapman & Hall, London 4th Edition
- 360 Hill, E.G., C.L. Silbernack, and W.O. Lundberg. 1965
Hypercholesterolemic effect of menhaden oil in the presence of dietary cholesterol in swine
Proc. Soc. Exp. Biol. Med. 119:368-370
- 361 Hill, E.G., E.L. Warmanen, C.L. Silbernack, and R.T. Holman. 1961
Essential fatty acid nutrition in swine. I. Linoleate requirement estimated from triene:tetraene ratio of tissue lipids
J. Nutr. 74:335

- 362 Hixson, A.W., and J.B. Bockelmann. 1942
Liquid-liquid extraction employing solvents in
the region of their critical temperatures
Trans. Am. Inst. Chem. Eng. 38:891-930
- 63 Hofmann, A.F. 1962
Separation of 1- and 2-monoglycerides by thin-
layer adsorption chromatography on hydroxyl
apatite
J. Lipid Res. 3:391-392
- 364 Hofmann, A.F. 1964
Hydroxyapatite as an adsorbent for thin-layer
chromatography: Separations of lipids and
proteins
In, New Biochemical Separations, A.T. James and
L.J. Morris (Ed.), D. Van Nostrand Co., London
- 365 Hofmann, Werner. 1937
Fetteinsparung bei fettgebacken. Gespritztes
teegebäck
Mehl Brot 37(33):3-4
- 366 Hofstetter, H.H., N. Sen, and R.T. Holman. 1965
Characterization of unsaturated fatty acids by
gas-liquid chromatography
J. Am. Oil Chem. Soc. 42:537-540
- 367 Holman, R. 1960
The ratio of trienoic: tetraenoic acids in tissue
lipids as a measure of essential fatty acid
requirement
J. Nutr. 70:405-410
- 368 Holman, R.T. 1954
Autoxidation of fats and related substances
In, Progress in the Chemistry of Fats and Other
Lipids, R.T. Holman, W.O. Lundberg, and T.
Malkin (Eds.), Acad. Press, N.Y. Vol. 2
- 369 Holman, R.T. 1956
Function and metabolism of essential fatty acids
In, Biochemical Problems of Lipids, Proceedings
of the Second International Conference, Univ. of
Ghent, July, 1955, G. Popjak and E. Le Breton
(Eds.), Butterworths Sci. Pub., London p. 463
- 370 Holman, R.T. 1957
Measurement of polyunsaturated fatty acids
In, Methods of Biochemical Analysis, D. Glick
(Ed.), Interscience Publishers, Inc., N.Y.
- 371 Holman, R.T. 1960
The ratio of trienoic tetraenoic acids in tissue
lipids as a measure of essential fatty acid
requirement
J. Nutr. 70:405
- * 372 Holman, R.T. 1964
Nutritional and metabolic interrelationships
between fatty acids
Fed. Proc. 23:1062-1067
- 373 Holman, R.T., and H. Hayes. 1958
Determination of polyunsaturated acids in lipids
of plasma and tissue
Anal. Chem. 30:1422
- 374 Holman, R.T., W.O. Lundberg, and T. Malkin (Eds.).
1952
Progress in the Chemistry of Fats and Other Lipids
Volume 1. Academic Press, New York. Volume 2.
1954. Academic Press, Inc., New York; Volume 3.
1955. Pergamon Press, New York; Volume 4. 1957.
Pergamon Press, New York; Volume 5. 1958.
Pergamon Press, New York; Volume 6. 1963. The
Macmillan Co., New York.
- 375 Holmboe, Carl Fred. 1938
Die energieumsetzung der katalytischen reaktion
von fetten olen mit wasserstoff
Ber. Dtsch. Chem. Ges. 71:532-541
- 376 Horning, E.C., and W.J.A. Vandeheuvel. 1964
Gas chromatography in lipid investigations
J. Am. Oil Chem. Soc. 41:707-716
- 377 Horning, E.C., A. Karmen, and C.C. Sweeley. 1964
Gas chromatography of lipids.
In, Progress in the Chemistry of Fats and Other
Lipids, R.T. Holman (Ed.), MacMillan Co., New
York
- 378 Horning, E.C., E.M. Ahrens, Jr., S.R. Lipsky,
F.H. Mattson, J.P. Mead, D.A. Turner, and W.H.
Goldwater. 1964
Quantitative analysis of fatty acids by gas-
liquid chromatography
J. Lipid Res. 5:20-27
- 379 Houchin, O.B. 1942
Vitamin E and muscle degeneration in the hamster
Fed. Proc. 1:117-118
- * 380 Houle, C.B. 1967
Gas-liquid chromatography
In, Fish Oils, Stansby, M.E. (Ed), AVI Publishing
Co., Westport, Conn. pp. 52-62
- 381 Hove, E.L. 1955
Anti-vitamin E stress factors as related to lipid
peroxides
Am. J. Clin. Nutr. 3:328
- 382 Hove, E.L., and P.L. Harris. 1946
Covitamin studies. 5. The interrelation of alpha-
tocopherol and essential unsaturated fat acids
J. Nutr. 31:699-744
- 383 Hove, E.L., and P.L. Harris. 1947
Relative activity of the tocopherols in curing
muscular dystrophy in rabbits
J. Nutr. 33:95
- 384 Idler, D.R., and V.H.M. Fagerlund. 1955
Marine sterols. I. Isolation of 24-
methylenecholesterol from molluscs
J. Am. Chem. Soc. 77:4142
- 385 Idler, D.R., T. Tamura, and T. Wainai. 1964
Seasonal variations in the sterol, fat and
unsaponifiable components of scallop muscle
J. Fish. Res. Board Can. 21:1035
- * 386 Idyll, C.P. 1973
The anchovy crisis
Sci. Am. 228(6):22-29
- 387 Imaichi, K., G.D. Michaels, B. Gunning, S.
Grasso, G. Fukayama, and L.W. Kinsell. 1963
Studies with the use of fish oil fractions in
human subjects
Am. J. Clin. Nutr. 13:158-168
- * 388 Imaichi, K., K. Oyama, J. Fukuda, M. Yamazaki,
and T. Hirose. 1965
Fatty acid composition of depot fat, in relation
to high fish consumption in Japan
Jap. Circ. J. 29:931-936
- 389 Ingold, H.K. 1962
Metal catalysis
In, Lipids and Their Oxidation, H.W. Schultz,
E.A. Day, and R.O. Sinnhuber (Eds.), AVI
Publishing Co., Westport, Conn.
- 390 Isselbacher, K.J. 1965
Metabolism and transport of lipid by intestinal
mucosa
Fed. Proc. 24:16-22
- 391 Ito, S., and K. Fukuzumi. 1963
Isolation and structure of eicosanoic,
docosanoic, octadecatetraenoic,
eicosapentaenoic, and docosahexaenoic acid in
skipper oil
J. Jap. Oil Chem. Soc. 12:272-277
- 392 Ito, S., and K. Fukuzumi. 1963
The composition of the component fatty acids of
several fish oils
J. Jap. Oil Chem. Soc. 12:278-281.

- 393 Iverson, J.L. 1965
Fatty acid content of vegetable oils, unusual oils, marine oils and margarines
J. Am. Oil Chem. Soc. 48:902-904
- 394 Iverson, J.L., D. Firestone, and J. Eisner. 1965
Evaluation of methods for identifying fatty acid esters separated by gas chromatography
J. Assoc. Offic. Agric. Chem. 48:482
- 395 Izumi, G. 1957
New uses for ozonized fats and oils. Production of surface active agents and plasticizers
Kogyo Kagaku Zasshi 60:730-734
- 396 IUPAC. 1966
Standard methods of the oils and facts section of the IUPAC
5th Ed., Butterworths, London (incorporating first supplement up-to-date 1965)
- 397 IUPAC. 1966
Impurities
IUPAC Standard Methods for the Analysis of Oils, Fats and Soaps II.C.2
- 398 IUPAC. 1966
Acid value
IUPAC Standard Methods for the analysis of Oils, Fats and Soaps 5th Ed., II.D.1.2
- 399 IUPAC. 1966
Peroxide value
IUPAC Standard Methods for the Analysis of Oils, Fats and Soaps 5th Ed., II.D.13
- 400 IUPAC. 1966
Moisture and volatile matter
IUPAC Standard Methods for the Analysis of Oils, Fats and Soaps 5th Ed., II.C.1.1
- 401 Jangaard, P.M. and R.G. Ackman. 1965
Lipids and component fatty acids of the Newfoundland squid (*Illex illecebrosus*)
J. Fish. Res. Bd. Can. 22:131-137
- 402 Jangaard, P.M. 1965
A rapid method for concentrating highly unsaturated fatty acid methyl esters in marine lipids as an aid in their identification by GLC
J. Am. Oil Chem. Soc. 42:845-847
- 403 Jantzen, E., and H. Andreas. 1961
Reaction of unsaturated fatty acids with mercury(II)-acetate. The use for preparative separation (in German)
Chem. Ber. 94:628-633
- 404 Jantzen, E., and O. Wieckhorst. 1954
Concentric tube columns for efficient distillation at low pressure (in German)
Chem. Ing. Tech. 26:392-396
- 405 Jantzen, E., H. Andreas, K. Morgenstern, and W. Roth. 1961
Separation of unsaturated fatty acids with the help of mercury adducts
Fette-Seifen-Anstrichm. 63:685-688
- 406 Jart, A. 1965
The fatty acid composition of some herring and whiting oils
Oleagineux 20 (7):447-448
- 407 Jessen, K.E., J. Glavind, S. Hartmann, and H. Dam. 1951
Peroxidation of human adipose tissue in peripheral venous diseases
Acta Path. Microbiol. Scand. 29:73-76
- 408 Johnson, K.L. 1964
Non ionic detergents derived from epoxidized oils
J. Am. Oil Chem. Soc. 41:191-198
- 409 Johnson, O.C. 1956
Chemical and nutritive changes in thermally oxidized edible fats and oils
Diss. Abstr. 16:1657
- 410 Johnson, O.C., and P.A. Kummerow. 1957
Chemical changes which take place in an edible oil during thermal oxidation
J. Am. Oil Chem. Soc. 34:407-409
- 411 Johnson, O.C., E. Perkins, M. Sugai, and P.A. Kummerow. 1957
Studies on the nutritional and physiological effects of thermally-oxidized oils
J. Am. Oil Chem. Soc. 34:594-597
- 412 Johnson, O.C., T. Sakuragi, and P.A. Kummerow. 1956
A comparative study of the nutritive value of thermally oxidized oils
J. Am. Oil Chem. Soc. 33:433-435
- 413 June, F.C. 1963
The menhaden fishery
In, Industrial Fishery Technology, H.E. Stansby (Ed.), Reinhold, N.Y. pp. 146-159
- 414 Jungermann, E., and P.E. Spoerri. 1953
The separation and identification of the 9,10- and 10,9-chlorohydrins of methyl oleate
J. Am. Chem. Soc. 75:4704-4705
- 415 Jungermann, E., and P.E. Spoerri. 1958
The reactions of hydrogen bromide with oleic acid and its esters. II. Free radical addition
J. Am. Oil Chem. Soc. 35:393-396
- 416 Kahn, S.G. 1964
A study of the hypercholesteremic activity of the ethyl esters of the polyunsaturated fatty acids of cod-liver oil in the rat
J. Nutr. 83:262-266
- 417 Kahn, S.G., J. Vandeputte, S. Wind, and H. Yacowitz. 1963
A study of the hypocholesterolemic activity of the ethyl esters of the polyunsaturated fatty acids of cod-liver oil in the chicken. I. Effect on serum cholesterol
J. Nutr. 80:403-413
- 418 Kahn, S.G., S. Wind, A. Slocum, D. Pfeffer, and H. Yacowitz. 1963
A study of the hypocholesterolemic activity of the ethyl esters of the polyunsaturated fatty acids of cod liver oil in the chicken. II. Effect on serum and tissue cholesterol and aortic and coronary atherosclerosis
J. Nutr. 80:414-424
- 419 Kamath, G.G., and N.G. Magar. 1956
Fatty acids and glycerides of shark (*Pristis*) liver oil
J. Indian Chem. Soc., Ind. News Ed. 19:171-176
- 420 Kamath, G.G., and N.G. Magar. 1956
Fatty acids and glycerides of shark (*Galeocerdo rayneri*) liver oil
J. Indian Chem. Soc., Ind. News Ed. 19:201-205
- 421 Kaneda, T., and R.B. Alfin-Slater. 1963
A Comparison of the effects of the polyunsaturated fatty acids of cuttle fish liver oil and cottonseed oil on cholesterol metabolism
J. Am. Oil Chem. Soc. 40:336-338
- 422 Kaneda, T., and S. Ishii. 1953
Studies on the nutritive value of lipids. VII. Nutritive value or toxicity of highly unsaturated fatty acids
Bull. Jap. Soc. Sci. Fish. 19:171-177
- 423 Kaneda, T., and S. Sakurai. 1954
Studies on the nutritive value of lipids. X. Seborrhea forming activity of esters of fatty acids with higher alcohols upon the rats (1)
Bull. Jap. Soc. Sci. Fish. 19:1168-1175
- 424 Kaneda, T., H. Sakai, S. Ishii, and K. Arai. 1955
Studies on the nutritive value of marine oils (In Japanese)
Bull. Tokai Reg. Fish. Res. Lab. No. 12

- 425 Kaneda, T., H. Sakurai, and S. Ishii. 1954
Studies on the nutritive value of lipids. XI.
Nutritive value or toxicity of
highly unsaturated fatty acids (2)
Bull. Jap. Soc. Sci. Fish. 20:50-57
- 426 Kaneda, T., H. Sakurai, and S. Ishii. 1954
Nutritive value or toxicity of highly unsaturated
fatty acids (3)
Bull. Jap. Soc. Sci. Fish. 20:658-663
- 427 Kaneda, T., S. Sakurai, and S. Ishii. 1957
Seborrhea forming activity of esters of fatty
acids with higher alcohols upon the rats (2)
Preventive methods of seborrhea
Bull. Jap. Soc. Sci. Fish. 23:324-330
- 428 Karmen, A., L. McCaffrey and R.L. Bowman. 1963
The validity of the results obtained when a
stream-splitting device is used before the gas-
liquid chromatography detector
J. Lipid Res. 4:108-110
- 429 Karnovsky, M.L., W.S. Rapson, and H.M. Schwartz.
1948
South African fish products. Part XXVIII. The
composition of the liver oil of the seven-gilled
shark, *Heptranchias pectorosus* (Ger.)
J. Soc. Chem. Ind. 67:144-147
- 430 Karnovsky, M.L., W.S. Rapson, and M. Black. 1946
South African fish products. Part XXIV. The
occurrence of alpha-glyceryl ethers in
the unsaponifiable fractions of natural fats
J. Soc. Chem. Ind. 65:425-428
- 431 Karrer, P., and H. Koenig. 1943
Essential unsaturated fatty acids
Helv. Chim. Acta 26:619-626
- 432 Karrick, N.L. 1965
A review of fish oil research conducted and
sponsored by the Bureau of Fish. of the U.S.
Fette, Seifen, Anstrichm. 67:489-494
- 433 Katada, M., K. Zama, and H. Igarashi. 1960
Lipids of the muscle of tuna (*Thynnus orientalis*).
VII. Distribution of the lipids in dark-colored
muscle, ordinary muscle, and several other tissues
Bull. Jap. Soc. Sci. Fish. 26:425-429
- 434 Kayama, M., Y. Tsuchiya, and J.F. Mead. 1963
A model experiment of aquatic food chain with
special significance in fatty acid conversion
Bull. Jap. Soc. Sci. Fish. 29:452-458
- 435 Kaufmann, H.P. and J. Baltes. 1936
Diene syntheses in the field of fats. I. The
diene numbers of fats
Fette, Seifen, Anstrichm. 43:93-97
- 436 Kaufmann, H.P. and T. Miyakawa. 1958
Marine fats. I. Fat content and fat constants
Fette, Seifen, Anstrichm. 60:469-482
- 437 Kaufmann, H.P. 1958
Analyse der fette und fettprodukte
Springer Verlag., Berlin
- 438 Kaufmann, H.P. 1962
Diels-Alder reactions in the field of fats
Fette-Seifen-Anstrichm. 64:1115-1126
- 439 Kaufmann, H.P., and E. Gottschalk. 1956
Body oil of the fish *Hoplostethus islandicus*
Fette Seifen Anstrichm. 58:411-446
- 440 Kaufmann, H.P., and H. Kirschnek. 1958
Fatty aldehydes. IV. Preparation of fatty
aldehydes containing more than one double bond
Fette, Seifen, Anstrichm. 60:1125-1132
- 441 Kaufmann, H.P., and H. Wessels. 1964
The thin-layer chromatography of fat bodies. XIV.
The separation of triglycerides by combinations
of adsorption and of reverse phase chromatography
(in German)
Fette, Seifen, Anstrichm. 66:81-86
- 442 Kaufmann, H.P., and T.H. Khoe. 1964
The analysis of fatty acids and triglycerides of
cod liver oil
Fette, Seifen, Anstrichm. 66:590-597
- 443 Kaunitz, H. 1960
Biological effects of rancid fats and their
functions
Exp. Med. Surg. 18:59
- 444 Kaunitz, H. 1962
Biological effects of autoxidized lipids
In, Symposium on Foods: Lipids and Their
Oxidation, H.W. Schultz (Ed.), AVI Publishing
Co., Westport, Conn.
- 445 Kaunitz, H., C.A. Slanetz, R.E. Johnson, H.B.
Knight, D.H. Saunders, and D. Swern. 1956
Biological effects of the polymeric residues
isolated from autoxidized fats
J. Am. Oil Chem. Soc. 33:630-634
- 446 Kaunitz, H., C.A. Slanetz, R.E. Johnson, J.
Guilmain, H.B. Knight, D.H. Saunders, and D.
Swern. 1956
Nutritional properties of the molecularly
distilled fractions of autoxidized fats
J. Nutr. 60:237-244
- 447 Kawai, S. 1942
Ester interchange between fatty oils and glycerol
J. Soc. Chem. Ind. Jap. 45:546-547
- 448 Kayama, M. 1964
Lipid metabolism in fish
J. Jap. Oil Chem. Soc. 13:511-519
- 449 Kayama, M., and Y. Tsuchiya. 1959
Intestinal absorption and distribution study of
oil in the carp, *Cyprinus carpio* Linnae
Tohoku J. Agric. Res. 10:229-236
- 450 Kayama, M., and Y. Tsuchiya. 1962
Possible conversion pathway of polyunsaturated
acid in fish
Tohoku J. Agric. Res. 13:229-235
- 451 Kayama, M., Y. Tsuchiya, and J.F. Mead. 1963
A model experiment of aquatic food chain with
special significance in fatty acid conversion
Bull. Jap. Soc. Sci. Fish. 29:452-458
- 452 Kayama, M., Y. Tsuchiya, J.C. Nevenzel, A. Fulco,
and J.F. Mead. 1963
Incorporation of linolenic-1-C14 acid into
eicosapentaenoic and docosahexaenoic acid in fish
J. Am. Oil Chem. Soc. 40:499-502
- 453 Kelly, P.B., R. Reiser, and D.W. Hood. 1958
The origin and metabolism of marine fatty acids;
the effect of diet on the depot fats of Mugil
cephalus (The Common Mullet)
J. Am. Oil Chem. Soc. 35:189-194
- 454 Kelly, P.B., R. Reiser, and D.W. Hood. 1958
The effect of diet on fatty acid composition of
several species of fresh water fish
J. Am. Oil Chem. Soc. 35:503-505
- 455 Kelly, P.B., R. Reiser, and D.W. Hood. 1959
The origin of the marine polyunsaturated fatty
acids. Composition of some marine plankton
J. Am. Oil Chem. Soc. 36:104-106
- 456 Kenney, H.E., D. Komanowsky, and A.N. Wrigley.
1965
Reaction of dichlorocarbene with fats and oils
J. Am. Oil Chem. Soc. 42:19-22
- 457 Kenney, H.E., D. Komanowsky, L.L. Cook, and A.N.
Wrigley. 1964
Preparation and etherification reaction of fatty
dichlorocyclopropanes
J. Am. Oil Chem. Soc. 41:82-85
- * 458 Keppler, J.G., J.A. Schols, W.H. Feenstra, and
P.W. Meijboom. 1963
Components of the hardening flavor present in
hardened linseed oil and soybean oil
J. Am. Oil Chem. Soc. 42:246-249

- 459 Keys, A., A. Kusukawa, B. Bronte-Stewart, N. Larsen, and M.H. Keys. 1958
Lessons from cholesterol studies in Japan, Hawaii, and Los Angeles
Ann. Intern. Med. 48:83-94
- 460 Kidger, D.P. 1968
Chocolate coating compositions
Can. Pat. 800,561 issued in 1968
- 461 King, A.E., H.L. Roschen, and W.H. Irwin. 1933
An accelerated stability test with the peroxide value as an index
Oil Soap 10:105-109
- 462 King, F.J., M.L. Anderson, and H.A. Steinberg. 1962
The effect of linoleic and linolenic acids on the solubility of cod actomyosin
J. Food Sci. 27:363-366
- 463 Kingsbury, K.J., G.M. Morgan, C. Aylott, and R. Emerson. 1961
Effects of ethyl arachidonate, cod liver oil, and corn oil on the plasma-cholesterol level
Lancet 1:739-741
- 464 Kinumaki, T., and K. Tabei. 1954-1955
Stabilizing effect of commercial antioxidants on vitamin A concentrates
Bull. Jap. Soc. Sci. Fish. 20:539-550
- 465 Kirschenbauer, H.G. 1960
Fats and oils, an outline of their chemistry and technology
Reinhold Publishing Corp., N.Y.
- 466 Kishimoto, T. 1966
Gangliosides in multiple sclerosis white matter, abstracted
Neurology 16:310
- 467 Kishimoto, Y., and M.S. Radin. 1964
Structures of the ester-linked mono- and diunsaturated fatty acids of pig brain
J. Lipid Res. 5:98-102
- * 468 Kishimoto, Y., M.S. Radin, W.W. Tourtellotte, J.A. Parker, and H.H. Itabashi. 1967
Gangliosides and glycerophospholipids in multiple sclerosis white matter
Arch. Neurol. 16:44-54
- 469 Klenk, E. 1957
Polyenoic acids of fish oils
In, Essential Fatty Acids, H.M. Sinclair (Ed.), Acad. Press, Inc. p. 57
- 470 Klenk, E. 1958
The polyenoic acids of fish oils
In, Essential Fatty Acids, Sinclair (Ed.), Butterworths, London pp. 57-59
- 471 Klenk, E. 1965
The metabolism of polyenoic fatty acids
J. Am. Oil Chem. Soc. 42:580-582
- 472 Klenk, E., and D. Eberhagen. 1962
About the composition of the fatty acid mixture of various fish oils
Z. Physiol. Chem. 328:180-188
- 473 Klenk, E., and D. Eberhagen. 1962
Unsaturated C16 fatty acids of marine plankton and the occurrence of delta,6,9,12,15-hexadecatetraenoic acid
Z. Physiol. Chem. 328:189-197
- 474 Klenk, E., and D. Eberhagen. 1957
The occurrence of delta-5,8,11,14,17-n-eicosapentaenoic acid in cod liver oil and its isolation
Z. Physiol. Chem. 307:42-48
- 475 Klenk, E., and G. Kremer. 1960
Biogenesis of C20 and C22 polyenoic acids in vertebrates
Z. Physiol. Chem. 320:111-125
- 476 Klenk, E., and H. Brockerhoff. 1957
The occurrence of delta-6,9,12,15-octadecatetraenoic acid in herring oil and its isolation
Z. Physiol. Chem. 307:272-277
- 477 Klenk, E., and H. Brockerhoff. 1958
The C18 and C22 polyenoic acids in herring oil
Z. Physiol. Chem. 310:153-170
- 478 Klenk, E., and H. Mohrhauer. 1960
Untersuchungen über den stoffwechsel der polyenfettsäuren bei der ratte
Z. Physiol. Chem. 320:218-232
- 479 Klenk, E., and H. Pfluger. 1963
The biogenesis of delta-5,8,11-eicosatrienoic acid in rats on a fat-free diet
Physiol. Chem. 335:53-62
- 480 Klenk, E., and H. Steinbach. 1959
The C16 polyenoic acids from herring oil
Z. Physiol. Chem. 316:31-44
- 481 Klenk, E., and H. Steinbach. 1959
The C16 polyenoic acids from herring oil
Z. Physiol. Chem. 320:111-125
- 482 Klenk, E., and K. Oette. 1960
The nature of C20 and C22 polyunsaturates occurring in the liver oils during the metabolism of linoleic acids by rats on a fat-free diet
Z. Physiol. Chem. 318:86-99
- 483 Klenk, E., and L. Brucker-Voight. 1961
The eicosapolyenoic acids of herring oil
Z. Physiol. Chem. 324:1-11
- 484 Klose, A.A., E.P. Mecchi, G.A. Behman, H. Lineveaver, F.H. Kratzer, and D. Williams. 1952
Chemical characteristics of turkey carcass fat as a function of dietary fat
Poult. Sci. 31:354-359
- 485 Klose, A.A., E.P. Mecchi, H.L. Hanson, and H. Lineveaver. 1951
The role of dietary fat in the quality of fresh and frozen storage turkeys
J. Am. Oil Chem. Soc. 28:162-164
- 486 Klose, A.A., H.L. Hanson, E.P. Mecchi, J.H. Anderson, I.V. Streeter, and H. Lineveaver. 1953
Quality and stability of turkeys as a function of dietary fat
Poult. Sci. 32:82-88
- 487 Knight, H.B., R.E. Koos, and D. Swern. 1954
New method for hydroxylating long-chain unsaturated fatty acids, esters, alcohols, and hydrocarbons
J. Am. Oil Chem. Soc. 31:1-5
- 488 Knipprath, W.G., and J.L. Mead. 1965
Influence of temperature on the fatty acid pattern of muscle and organ lipids of the rainbow trout (*Salmo gairdneri*)
Fish. Ind. Res. 3(1):23-27
- 489 Konen, J.C., E.T. Clocker, and R.P. Cox. 1945
Esterification of unsaturated fatty acids
Oil Soap 22:57-60
- 490 Kooy, H.J. 1964
Refining process for fish oils
Dutch Pat. 260,201 issued Jan. 27, 1964
- * 491 Koritala, S. 1968
Selective hydrogenation of soybean oil. III. Copper-exchanged molecular sieves and other supported catalysts
J. Am. Oil Chem. Soc. 45:197-200
- 492 Kornerup, V. 1950
Concentration of cholesterol, total fat, and phospholipids in serum of normal man
A.M.A. Arch. Intern. Med. 85:398-415

- 493 Kritchevsky, D., and S.A. Tepper. 1961
The free and ester sterol content of various
foodstuffs
J. Nutr. 74:441-444
- 494 Kritchevsky, D., A. Moyer, W.C. Tassar, R.F.J.
MacCandles, J.B. Logan, R.A. Brown, and M.E.
Englert. 1956
Cholesterol vehicle in experimental
atherosclerosis. II. Influence of unsaturation
Am. J. Physiol. 135:279-280
- 495 Kritchevsky, D., A.W. Mayer, W.C. Tassar, J.B.
Logan, R.A. Brown, and G. Richmond. 1954
Squalene feeding in experimental atherosclerosis
Circ. Res. 2:340-343
- * 496 Kritchevsky, D., S.A. Tepper, M.W. DiTullo, and
W.L. Holmes. 1967
The sterols of seafood
J. Food Sci. 32:64-66
- 497 Laksesvela, B. 1966
Sildefett som kyllingfett. (Herring fat as chicken
feed)
Rep. Herring Oil and Herring Meal Industry,
Bergen. Meld. SSF 4:92-102
- 498 Lalor, R.J., W.L. Leoschke, and C.A. Elvehjem.
1951
Yellow fat in the mink
J. Nutr. 45:183-188
- 499 Lambertsen, G., and O.R. Braekkan. 1965
Reports on technological research concerning
Norwegian fish industry
Bergen, Norway 4(13)
- 500 Lambertsen, G., and O.R. Braekkan. 1965
The fatty acid composition of herring oils
Fisk. Dir. Skr. (Ser. Teknol. Undersok.)
4(13):1-15
- 501 Lambertsen, G., and R.T. Holman. 1963
Partial characterization of the hydrocarbons of
herring oil
Acta Chem. Scand. 17(4): 281-282
- * 502 Lambertsen, G., H. Myklestad and O.R. Braekkan.
1971
Studies on monoene fatty acid isomers in
hydrogenated fish oils
J. Am. Oil Chem. Soc. 48:389-391
- * 503 Lambertsen, G., H. Myklestad, and O.R. Braekkan.
1966
The fatty acid composition of commercially
hydrogenated fish oils
J. Food Sci. 31:48-52
- 504 Lambertsen, G., H. Myklestad, and O.R. Braekkan.
1964
The determination and contents of alpha- and
gamma-tocopherols in margarine
J. Food Sci. 29:164-167
- 505 Lane, I.W., and H.H. Kriegel. 1963
The story of menhaden fish oil
J. Am. Oil Chem. Soc. 40(4-5):34
- 506 Larsen, N.P., and W.M. Bortz. 1959
Atherosclerosis: a comparative study of
Caucasian and Japanese citizens in the Hawaiian
Islands
J. Am. Geriatr. Soc. 8:867-872
- 507 Lassen, S. 1963
The sardine, mackerel, and herring fisheries
In, Industrial Fishery Technology, M.E. Stansby
(Ed.), Reinhold Publishing Corporation, N.Y.
10:131-145
- 508 Lassen, S., E.K. Bacon, and H.J. Dunn. 1949
The digestibility of polymerized oils
Arch. Biochem. 23:1-7
- 509 Le Roux, B.D.C. 1962
Butylated hydroxytoluene as antioxidant in fish
meal
Fish. Ind. Res. Inst., Cape Town, S. Afr. Annu.
Rep. pp. 58-60
- 510 Lea, C.H. 1931
The effect of light on the oxidation of fats
Proc. Roy. Soc. B. 108:176-189
- 511 Lea, C.H. 1936
Influence of cod liver oil in the diet on
susceptibility to oxidation of fat of pig
Dept. Sci. Ind. Res., Rep. Food Invest. Board.
pp. 73-75
- * 512 Lea, C.H. 1936
Antioxidants and the preservation of edible fats
J. Soc. Chem. Ind. (Trans. & Comm.) 55:293-302
- 513 Lea, C.H. 1939
Rancidity in Edible Fats
Chemical Publishing Co., N.Y.
- 514 Lea, C.H. 1953
Recent developments in the study of oxidative
deterioration of lipids
Chem. Ind. p. 1303
- 515 Lea, C.H. 1965
Chemical and nutritive aspects of oxidized and
heated fats
Chem. Ind. 6:244-248
- 516 Lea, C.H., L.J. Parr, and K.J. Carpenter. 1958
Chemical and nutritional changes in stored
herring meal
Brit. J. Nutr. 12:297-312
- 517 Lea, C.H., L.J. Parr, and K.J. Carpenter. 1960
Chemical and nutritional changes in stored
herring meal. 2
Brit. J. Nutr. 14:91-113
- 518 Lea, C.H., L.J. Parr, J.L. L'Estrange, and K.J.
Carpenter. 1966
Nutritional effects of autoxidized fats in animal
diets. 3. The growth of turkeys on diets
containing autoxidized fish oil
Brit. J. Nutr. 20:123-133
- 519 Leach, T.A., and W.S. Golding. 1931
Preliminary report of substitution of pilchard
oil for butterfat in milk for calf feeding
Sci. Agr. 12:204-205
- 520 Lee, C.P. 1953
Menhaden industry - past and present
U.S. Dept. of the Interior, Fish and Wildlife
Service, Fish. Leaflet 412:17 pp.
- 521 Lee, C.P. 1963
Processing fish meal and oil
In, Industrial Fishery Technology, M.E. Stansby
(Ed.), Reinhold Publishing Corp., N.Y.
pp. 219-235
- 522 Lehman, L.W., and E.J. Gauglitz, Jr. 1966
The preparation of alkyl esters from highly
unsaturated triglycerides. II
J. Am. Oil Chem. Soc. 43:383-384
- 523 Leong, K.C., G.M. Knobl, Jr., D.G. Snyder, and
E.H. Gruger, Jr. 1964
Feeding of fish oil and ethyl ester fractions of
fish oil to broilers
Poult. Sci. 43:1235-1240
- 524 Levin, Ezra, and R.K. Finn. 1955
A process for dehydrating and defatting tissues
at low temperature
Chem. Eng. Prog. 51:223-226
- 525 Lewis, R.W. 1962
Temperature and pressure effects on the fatty
acids of some marine ectotherms
Comp. Biochem. Physiol. 6:75-89
- 526 Lillard, D.A., and E.A. Day. 1961
Autoxidation of milk lipids. II. The relationship
of sensory to chemical methods for measuring the
oxidized flavor of milk fats
J. Dairy Sci. 44:623-632

- 527 Liston, J., M.E. Stansby, and H.S. Olcott. 1963
Bacteriological and chemical basis for
deteriorative changes
In, *Ind. Fish. Tech.*, M.E. Stansby (Ed.),
Reinhold, N.Y. pp. 350-361
- 8 Longenecker, H.E. 1937
An efficient fractionation equipment for the
qualitative and quantitative examination of
natural fats
J. Soc. Chem. Ind. (London) 56:199T-202T
- 529 Lovern, J. 1965
Some analytical problems in the analysis of fish
and fish products
J. Assoc. Offic. Agric. Chem. 48:60-68
- 530 Lovern, J.A. 1935
Fat metabolism in fishes. VI. The fats of some
plankton crustacea
Biochem. J. 29:847-849
- 531 Lovern, J.A. 1937
Fat metabolism in fishes. XI. Specific
peculiarities in depot fat composition
Biochem. J. 31:755-763
- 532 Lovern, J.A. 1938
Fat metabolism in fishes. XII. Seasonal changes
in the composition of herring fat
Biochem. J. 32:676-680
- 533 Lovern, J.A. 1942
The composition of the depot fats of aquatic
animals
Food Investigation Board Special Report, Dept.
Scientific and Industrial Research, H.M.
Stationery Office, York House, Kingsway, London
No. 51
- 534 Lovern, J.A. 1951
The biochemistry and metabolism of fats in fish
In, *Biochem. Soc. Sym.*, Williams (Ed.), Cambridge
Univ. Press, London 6:49-62
- 335 Lovern, J.A. 1962
The lipids in fish and the changes occurring in
them during processing and cold storage
In, *Fish in Nutrition*, Heen and Dreuzer (Eds.),
Fishing News, London pp. 86-111
- 536 Lovern, J.A. 1964
The lipids of marine organisms
Oceanogr. Mar. Biol. Ann. Rev. 2:169-191
- 537 Lucentini, Renato, and Elsa Drago. 1935
Eine neue Butterzahl
Ann. Chim. Appl. 25:388-391
- 538 Ludorff, W. 1937
Heringssöl: ein Beitrag zur schliessung der
fettlucke
Fette Seifen 44:416-420
- 539 Lundberg, W.O. 1949
Oxidation of esters of linoleic acid by oxygen
Oleagineux 4:86-94
- 540 Lundberg, W.O. 1961, 1962
Autoxidation and Antioxidants
Interscience Publishers, N.Y. I & II
- * 541 Lunde, G. 1972
Analysis of arsenic and bromine in marine and
terrestrial oils
J. Am. Oil Chem. Soc. 49:44-47
- 542 Lyles, C.H. 1966
Fisheries of the U.S. - 1965
U.S. Fish and Wildlife Service, C.F.S. (Current
Fishery Statistics) 4100:65 pp.
- 543 Lynen, F. 1961
Biosynthesis of fatty acids
In, *Drugs Affecting Lipid Metabolism*, Garattini
and Paoletti (Eds.), Elsevier Publ., Amsterdam
- 544 Lyness, W.I., and F.W. Quackenbush. 1955
Addition of chlorine to unsaturated fatty acids
and esters
J. Am. Oil Chem. Soc. 32:520-521
- 545 Machlin, L.J. 1962
Effect of dietary linolenate on the proportion of
linoleate and arachidonate in liver fat
Nature, London 194:868-869
- 546 Machlin, L.J. 1962
Effect of dietary linolenate on the proportion of
linoleate and archidonate in liver fat
Nature 194:868
- 547 Machlin, L.J., and R.S. Gordon. 1960
Linoleic acid as causative agent of
encephalomalacia in chickens fed oxidized fats
Proc. Soc. Exp. Biol. Med. 103:659
- 548 Machlin, L.J., R.S. Gordon, J.E. Marr, and C.W.
Pope. 1962
Effect of antioxidants and unsaturated fatty
acids on reproduction of the hen
J. Nutr. 76:284
- 549 Mack, C.H., and W.G. Bickford. 1953
Hydroxystearic acids. I. The catalytic
hydrogenation of the 9,10-epoxystearates
J. Org. Chem. 18:686-692
- 550 Mackenzie, C.G., and E.V. McCollum. 1940
The cure of nutritional muscular dystrophy in the
rabbit by alpha-tocopherol and its effect on
creatine metabolism
J. Nutr. 19:345
- 551 Mackenzie, J.B., and C.G. Mackenzie. 1953
Vitamin E activity of alpha-
tocopherylhydroquinone and muscular dystrophy
Proc. Soc. Exp. Biol. 84:388-392
- 552 MacKenzie, C.G., J.B. MacKenzie, and E.V. McCollum
1941
Uncomplicated vitamin E deficiency in the rabbit
and its relation to the toxicity of cod liver
oil
J. Nutr. 21:225
- 553 Madsen, L.L. 1936
Comparative effects of cod liver oil, cod liver
oil concentrate, lard, and cottonseed oil in a
synthetic diet on development of nutritional
muscular dystrophy
J. Nutr. 11:471-489
- 554 Madsen, L.L., C.M. McCay, and L.A. Maynard. 1933
Possible relationship between cod liver oil and
muscular degeneration of herbivora fed synthetic
diets
Proc. Soc. Exp. Biol. Med. 30:1434-1438
- 555 Madsen, L.L., C.M. McCay, and L.A. Maynard. 1935
Synthetic diets for herbivora, with special
reference to the toxicity of cod-liver oil
Cornell Univ. Agric. Exp. Sta. Memoir. p. 178
- 556 Maier, R., and H.K. Mangold. 1964
Thin-layer chromatography
In, *Advances in Analytical Chemistry and
Instrumentation*, C.N. Reilly (Ed.)
Interscience Publishers, John Wiley and Sons,
N.Y.
- 557 Malins, D.C. 1960
Fatty acids and glyceryl ethers in alkoxy-
diglycerides of dogfish liver oil
Chem. Ind. 44:1359-1360
- * 558 Malins, D.C. 1967
The classes of lipids in fish
In, *Fish Oils*, Stansby, M.E. (Ed.), AVI
Publishing Co., Westport, Conn. pp. 31-42
- 559 Malins, D.C., and C.R. Houle. 1961
Monoenoic fatty acids in dogfish livers: isomers
of the C16, C18, C20 and C22 series
Proc. Soc. Exp. Biol. Med. 108:126-129

- * 560 Malins, D.C., and H.K. Mangold. 1960
Analysis of complex lipid mixtures by thin-layer chromatography and complementary methods
J. Am. Oil Chem. Soc. 37:576-578
- 61 Malins, D.C., and J.C. Wekell. 1967
Chromatographic methods
In, Fish Oils, Stansby, M.E. (Ed.), AVI Publishing Co., Westport, Conn. pp. 63-72
- 562 Malins, D.C., J.C. Wekell, and C.R. Houle. 1965
Composition of the diacyl glyceryl ethers and triglycerides of the flesh and liver of the dogfish (*Squalus acanthias*)
J. Lipid Res. 6:100-105
- 563 Mangold, H.K. 1959
Synthesis of unsaturated fatty aldehydes
J. Org. Chem. 24:405-407
- * 564 Mangold, H.K. 1961
Thin layer chromatography of lipids
J. Am. Oil Chem. Soc. 38:708-727
- 565 Mangold, H.K., and D.C. Malins. 1960
Fractionation of fats, oils, and waxes on thin layers of silicic acid
J. Am. Oil Chem. Soc. 37:383-385
- 566 Mangold, H.K., and R. Kammerer. 1961
Separation, identification, and quantitative analysis of fatty acids by thin-layer chromatography and gas-liquid chromatography
Chem. Ind. (London) pp. 1032-1034
- 567 Maplesden, D.C., and J.K. Loosli. 1960
Nutritional muscular dystrophy in calves. II. Addition of selenium and tocopherol to a basal dystrophogenic diet containing cod liver oil
J. Dairy Sci. 43:645-653
- 568 Marble, D.R., J.E. Hunter, H.C. Knandel, and R.A. Dutcher. 1938
Fishy flavor and odor in turkey meat
Poult. Sci. 17:49-53
- 569 March, B.E., J. Biely, F.E. Claggett and H.L.A. Tarr. 1962
Nutritional and chemical changes in the lipid fraction of herring meals with and without antioxidant treatment
Poult. Sci. 41:873-880
- 570 Marion, J.B., and H.M. Edwards, Jr. 1964
Response of laying hens to dietary oils and purified fatty acids
Poult. Sci. 43:911-918
- 571 Markley, K.S. 1960
Fatty Acids
Interscience Publishers, New York Part I
- 572 Markley, K.S. 1961
Fatty Acids
Interscience Publishers, N.Y. Part II
- 573 Markley, K.S. 1961
Hydrogenation
In, Fatty Acids, Their Chemistry, Properties, Production, and Uses, 2nd Ed., K.S. Markley (Ed.), Interscience Publishers, N.Y.
- 574 Markley, K.S. 1964
Salt solubility methods, low temperature crystallization
In, Fatty Acids, 2nd Ed., Interscience Publishers, N.Y. Part III:2072-2123
- 575 Markley, K.S. 1964
Distillation
In, Fatty Acids, 2nd Ed., Interscience Publishers, N.Y. Part III:2016-2071
- 576 Markley, K.S. 1964
Sulfur derivatives
In, Fatty Acids, Their Chemistry, Properties, Production, and Uses, 2nd Ed., K.S. Markley (Ed.), Interscience Publishers, N.Y.
- 577 Martin, A.J.P., and T. Moore. 1939
Some effects of prolonged vitamin E deficiency in the rat
J. Hyg. Camb. 39:643-650
- 578 Mason, K.E. 1940
Minimal requirements of male and female rats for vitamin E
Am. J. Physiol. 134:268
- 579 Mason, K.E., and A.P. Emmel. 1945
Vitamin E and muscle pigment in the rat
Anat. Rec. 92:33-50
- 580 Mason, K.E., and G.R. Hartsough. 1951
Relation of "steatitis" or "yellow fat" in mink to dietary fats and inadequacy of vitamin E
Fed. Proc. 10:389
- 581 Mason, K.E., and G.R. Hartsough. 1951
"Steatitis" or "yellow fat" in mink and its relation to dietary fats and inadequacy of vitamin E
J. Am. Vet. Med. Assoc. 119:72
- 582 Mason, K.E., H. Dam, and H. Granados. 1946
Histological changes in adipose tissue of rats fed a vitamin E deficient diet high in cod liver oil
Anat. Rec. 94:265
- 583 Matic, M. 1958
South African pilchard oil. 7. The isolation and structure of an octadecatetraenoic acid from South African pilchard oil
Biochem. J. 68:692-695
- 584 Matsumoto, T., H. Sone, and I. Niiya. 1955
The lipid of a fish, *Xenogramma carinatum*
J. Jap. Oil Chem. Soc. 4:131-133
- * 585 Matsuo, N. 1954
Studies on the toxicity of fish oil. II
J. Biochem. 41(5):647-652
- * 586 Matsuo, N. 1954
Studies on the toxicity of fish oil
J. Biochem. (Japan) 41(4):481-487
- 587 Matsuo, N. 1957
Studies on the toxicity of fish oils. III
J. Jap. Biochem. Soc. 29:769-772
- 588 Matsuo, N. 1957
Studies on the toxicity of fish oils. IV. The effects of autoxidized ethyl ester of highly unsaturated fatty acid on protein and amino acid
J. Jap. Biochem. Soc. 29:773-777
- 589 Matsuo, N. 1957
Toxicity of fish oil
Kagaku No Ryoiki 12:64-73
- 590 Matsuo, N. 1957
Studies on the toxicity of fish oils. VII. Studies on seborrhea of rat
J. Jap. Biochem. Soc. 29:816-818
- 591 Matsuo, N. 1957
Studies on the toxicity of fish oils. VIII. Toxicity of fish oil polymerized by heating under carbon dioxide
J. Jap. Biochem. Soc. 29:885-891
- 592 Matsuo, N. 1958
Studies on the toxicity of fish oils. IX. Toxicity of fish oil heat-polymerized in the presence of the air
J. Jap. Soc. Food Nutr. 10:255-259
- 593 Matsuo, N. 1959
Biochemical studies on oils and fats. XIII. Toxicity of thermally polymerized ethyl linolenate
J. Jap. Soc. Food Nutr. 12:206-209

- 594 Matsuo, N. 1959
Biochemical studies on oils and fats. XIV.
Toxicity of cyclic compounds
J. Jap. Soc. Food Nutr. 12:210-213

Matsuo, N. 1960
Studies on the toxicity of fish oils. XI.
Toxicity of heat-polymerized oil
J. Jap. Oil Chem. Soc. 9:37-41
- 596 Matsuo, N. 1960
Biochemical studies on oils and fats. XII.
Toxicity of cyclic compound of beta-eleostearic acid
J. Chem. Soc. Jap., Pure Chem. Sect. 81:469-472
- * 597 Matsuo, N. 1961
Studies on the toxicity of fish oil (V). Toxicity of oxidized commercial cod-liver oil
Tokushima J. Exp. Med. 8:96-100
- * 598 Matsuo, N. 1961
Studies on the toxicity of fish oil (III). Toxicity of autoxidized highly unsaturated fatty acid ethyl ester applied to rat
Tokushima J. Exp. Med. 8:90-95
- * 599 Matsuo, N. 1962
Nutritional effects of oxidized and thermally polymerized fish oils
Symposium on Foods: Lipids and their Oxidation. AVI Publishing Co., Westport Conn. pp. 321-359
- 600 Mattei, V., and W.T. Roddy. 1959
The use of fish oils for fatliquoring leather. I. Menhaden oil and cod oil fatliquors. II. Use of phosphoric acid to control rate of fish oil sulfation. III. Suitability of ocean perch, herring, salmon, and menhaden oils in fatliquoring
J. Am. Leather Chem. Assoc. 54:12-24, 555-567, 640-653
- 601 Mattikow, M. 1942
A new method of refining oils with nonsaponifying alkalis-the Clayton process
Oil Soap 19:83-87
- 602 Mattil, K.F. 1964
Hydrogenation
In, Bailey's Industrial Oil and Fat Products, D. Sverner (Ed.), Interscience Publishers, N.Y.
- 603 Mattil, K.F. 1964
Deodorization
In, Bailey's Industrial Oil and Fat Products, D. Sverner (Ed.), Interscience Publishers, N.Y.
- 604 Mattill, H.A., and C. Golumbic. 1942
Vitamin E, cod liver oil, and muscular dystrophy
J. Nutr. 23:625-631
- 605 Maynard, L.A., C.M. McCay, and L.L. Madsen. 1936
Influence of food fat of varying degrees of unsaturation upon blood lipids and milk fat
J. Dairy Sci. 19:49-53
- 606 McArdle, E.H. 1931
Core oils for use in making foundry cores
U.S. Pat. 1,822,229 issued Sep. 8, 1931
- * 607 McCay, C.M. and H. Paul. 1938
The effect of melting point of fat upon its utilization by guinea pigs
J. Nutr. 15:377-382
- 608 McCay, C.M. 1937
The biochemistry of fish
Ann. Rev. Biochem. 6:445
- 609 McCay, C.M., and L.A. Maynard. 1935
Effect of ingested cod-liver oil, shark-liver oil, and salmon oil on composition of blood and milk of lactating cows
J. Biol. Chem. 109:29-37
- 610 McCay, C.M., H. Paul, and L.A. Maynard. 1938
Influence of hydrogenation and of yeast in counteracting cod-liver oil injury in herbivora and the influence of salmon oil on fat secretion
J. Nutr. 15:367-375
- 611 McCharty, R.C. 1964
Mammalian metabolism of straight-chain saturated hydrocarbons
Biochem. Biophys. Acta 84:74-79
- 612 McGrath, B.P., and J.M. Tedder. 1961
The mechanism of allylic bromination by N-bromosuccinimide
Proc. Chem. Soc. (London) 1961(Feb):80-81
- 613 McKay, A.F., and A.R. Bader. 1948
The preparation and properties of the eight diastereoisomers of 9,10,12,13-tetrahydroxystearic acid
J. Org. Chem. 13:75-85
- 614 Mead, J.D., and D.R. Howton. 1957
Metabolism of essential fatty acids. VII. Conversion of gamma-linolenic acid to arachidonic acid
J. Biol. Chem. 229:575-582
- 615 Mead, J.F. 1960
The metabolism of the unsaturated fatty acids
In, Lipid Metabolism, K. Block (Ed.), John Wiley & Sons, N.Y.
- 616 Mead, J.F., and M. Kayama. 1967
Lipid metabolism in fish
In, Fish Oils, M.E. Stansby (Ed.), AVI, Westport, Conn. pp. 289-299
- 617 Mead, J.F., M. Kayama, and R. Reiser. 1960
Biogenesis of polyunsaturated fatty acids in fish
J. Am. Oil Chem. Soc. 37:438-440
- 618 Mehlenbacher, V.C. 1960
The Analysis of Fats and Oils
The Garrard Press, Champaign, Ill. pp. 502-504
- 619 Mehlenbacher, V.C., T.H. Hopper, and E.M. Saltee. 1965
Official and tentative methods of the American Oil Chem. Soc.
Am. Oil Chem. Soc., N.Y.
- * 620 Meijboom, P.W. 1964
Relationship between molecular structure and flavor perceptibility of aliphatic aldehydes
J. Am. Oil Chem. Soc. 41:326-328
- * 621 Melnick, D., C.M. Gooding and H.W. Vahlteich (The Best Foods, Inc. New York). 1952
Soya Bean Oil Compositions
U.S. Patent 2,627,469
- * 622 Menzel, D.B., and H.S. Olcott. 1964
Positional distribution of fatty acids in fish and other animal lecithins
Biochim. Biophys. Acta 84:133-139
- 623 Mercier, J. 1962
Effect of irradiation on food fats
Ann. Nutr. Aliment. 16:59-89
- * 624 Merker, D.R. (Swift & Co., Chicago, Ill.). 1957
Deodorization of hydrogenated fatty oils
U.S. Patent 2,783,260, issued Feb. 26, 1957
- * 625 Merker, D.R. (Swift & Co., Chicago, Ill.). 1958
Non-selective hydrogenation of fats and oils
U.S. Patent 2,862,941, issued Dec. 2, 1958
- 626 Meszaros, G. 1935
Über die emulgierbarkeit der speisefette
Z. Unters. Lebensm. 69:318-330
- 627 Metcalfe, L.D., and A.A. Schmitz. 1961
The rapid preparation of fatty acid esters for gas chromatographic analysis
Anal. Chem. 33:363-364
- 628 Mezzalana, G. 1960
Sur un nouvel anticholestérolémiant obtenu par extraction de l'huile de poisson
Minerva Med. 51:646

- 629 Michaelis, L., and S.H. Wollman. 1949
The semiquinone radical of tocopherol
Science 109:313-314
- 630 Miller, D., K.C. Leong, G.M. Knobl, and E. Gruger. 1964
Exudative diathesis and muscular dystrophy induced in the chick by esters of polyunsaturated fatty acids
Proc. Soc. Exp. Biol. 116:1147-1151
- 631 Miller, S.A., A. Dymaza, and S.A. Goldblith. 1962
Cholesterolemia and cardiovascular sudanophilia in rats fed sardine mixtures
J. Nutr. 77:397-402
- 632 Mitchell, J., Jr., D.M. Smith, and W.M. Eryant. 1940
Analytical procedure employing Karl Fischer reagent. III. The determination of organic acids
J. Am. Chem. Soc. 62:4-6
- 633 Mitchell, J.H., H.R. Kraybill, and P.P. Zscheile. 1943
Quantitative spectral analysis of fats
Ind. Eng. Chem. Anal. Ed. 15:1-3
- 634 Miwa, T.K., K.L. Mikolajczak, F.R. Earle, and I.A. Wolff. 1960
Gas chromatographic characterization of fatty acids. Identification constants for mono- and dicarboxylic methyl esters
Anal. Chem. 32:1739-1742
- * 635 Miyake, R., Y. Shimamura, and H. Yajima. 1965
Kinetics of fat hydrogenation
Chem. Pharm. Bull. 13(7):843-858
- 636 Mohrhauer, H., and R.T. Holman. 1963
Effect of linolenic acid upon the metabolism of linoleic acid
J. Nutr. 81:67-74
- 537 Mohrhauer, H., and R.T. Holman. 1963
The effect of dose level of essential fatty acids upon fatty acid composition of the rat liver
J. Lipid Res. 4(2):151-159
- * 638 Mohrhauer, H., K. Christiansen, M.V. Gan, M. Deubig, and R.T. Holman. 1967
Chain elongation of linoleic acid and its inhibition by other fatty acids in vitro
J. Biol. Chem. 242:4507-4514
- 639 Monick, J.A., H.D. Allen, and C.J. Marlies. 1946
Vapor-liquid equilibrium data for fatty acids and fatty methyl esters at low pressures
J. Am. Oil Chem. Soc. 23:177-182
- 640 Moore, T. 1937
Spectroscopic changes in fatty acids. I. Changes in the absorption spectra of various fats induced by treatment with potassium hydroxide
Biochem. J. 31:138-141
- 641 Moore, T., and I.M. Sharman. 1961
Prevention of injurious effects of excessive cod-liver oil by its fortification with vitamin E
Brit. J. Nutr. 15:297-303
- 642 Moore, T., I.M. Sharman, and J.R. Ward. 1953
The vitamin E activity of methylene blue
Biochem. J. 56:xxx1
- 643 Moore, T., I.M. Sharman, and J.R. Ward. 1953
The vitamin E activity of substances related to methylene blue
Biochem. J. 54:xvi-xvii
- 644 Moore, T., I.M. Sharman, and R.J. Ward. 1953
The partial vitamin E activity of malachite green
Proc. Nutr. Soc. 12:2,v
- 645 Moore, T., I.M. Sharman, and R.J. Ward. 1954
The partial vitamin E activity of certain redox dyes
Biochem. J. 58:vii
- 646 Moore, T., I.M. Sharman, and R.J. Ward. 1959
Cod-liver oil as source and antagonist of vitamin E
Brit. J. Nutr. 13:100-110
- 647 Moreno, J.M.M., and A.V. Roncero. 1964
The application of urea inclusion compounds in fat analysis
In, Analysis and Characterization of Oils, Fats and Fat Products, H.A. Boekennoogen (Ed.), Interscience Publishers, N.Y. Vol. 1
- 648 Mork, P.C. 1972
Heterogeneous hydrogenation of fish oils: Kinetic determination of catalyst poisoning
Paper presented at the Eleventh World Congress, ISF, Goteborg, Sweden, June, 1972.
- * 649 Morris, C.E. and F.P. Khym. 1945
Recovery of catalyst from hydrogenated oil
U.S. Patent 2,602,807
- 650 Mukherjee, S. 1955
A rapid method for the estimation of unsaturation of fats and oils by use of hypochlorous acid reagent
J. Am. Oil Chem. Soc. 32:351-353
- 651 Munson, T.O., J. Holzworth, E. Small, S. Witzel, T.C. Jones, and H. Luginbuhl. 1958
Steatitis ("Yellow fat") in cats fed canned red tuna
J. Am. Vet. Med. Assoc. 133:563-568
- 652 Murphy, R.R., R.V. Boucher, and H.C. Kandel. 1939
Flavor of turkey meat as affected by feeding fish meal and oil
Proc. World's Poul. Congr. Expo., 7th Congr. pp. 542-545
- 653 Murray, K.E. 1955
Low pressure fractional distillation and its use in the investigation of lipids
In, Progress in the Chemistry of Fats, R.T. Holman, W.O. Lundberg, and T. Malkin (Eds.), Pergamon Press, N.Y.
- 654 Nakamura, M. 1937
Studies on peroxide of oil
J. Soc. Chem. Ind. Jap. 40:442-452
- 655 Nanavati, D.D., B. Nath, and J.S. Aggarwal. 1959
N-Bromosuccinimide in the conjugation of fatty acids and oils
J. Am. Oil Chem. Soc. 36:226-230
- 656 Nason, A., and I.R. Lehman. 1955
Tocopherol as an activator of cytochrome C reductase
Science 122:19-22
- * 657 National Academy of Sciences, Committee on Nutritional Misinformation. 1973
Supplementation of Human Diets with Vitamin E
A Statement of the Food & Nutrition Board, Div. of Biol. and Agr., National Research Council
- 658 Naudet, M., and E. Ucciani. 1960
About the bromination in the allylic position of oleic acid methyl ester
Fette-Seifen-Anstrichm. 62:691-694
- 659 Nelson, V.E., R.L. Jones, G. Adams, and L.T. Anderegg. 1927
Cod-liver oil as food-observations on the existence of vitamin E
Ind. Eng. Chem. 19:840-844
- * 660 Neudoerffer, T.S., and C.H. Lea. 1966
Effects of dietary fish oil on the composition and stability of turkey depot fat
Brit. J. Nutr. 20:581-594
- 661 Ni, T.C. 1938
Further experiments on the prevention of nutritional encephalomalacia in chickens
Chin. J. Physiol. 13:229

- 662 Ni, T.C. 1940
Nutritional encephalomalacia and some factors
accelerating its onset
Chin. J. Physiol. 15:181
- 663 Nicholaysen, R., and A. Pihl. 1953
Feeding experiments in rats and in man with a
commercial heat-treated herring oil
Tidsskr. Kjem. Bergvesen Met. 13:5-7
- 664 Nichols, P.L., Jr. 1952
Coordination of silver ion with methyl esters of
oleic and elaidic acids
J. Am. Chem. Soc. 74:1091-1092
- 665 Nicolaides, N., and A.N. Woodall. 1962
Impaired pigmentation in chinook salmon fed diets
deficient in essential fatty acids
J. Nutr. 78:431-437
- 666 Nicolet, B.H., and C.H. De Milt. 1927
A phenylstearic acid from oleic acid
J. Am. Chem. Soc. 49:1103-1106
- 667 Nieto, M. 1962
Technology of anchovy oil
Rev. Ing. Quim Ind. 4
- * 668 Njaa, L.R., O.R. Braekkan, G. Lambertsen, and P.
Utne. 1971
Hydrogenated marine fat, effect on growth and
liver weight in young rats given a high fat, low
protein ration
Nutr. Metabolism 13:207-221
- 669 Nonaka, J. 1957
Studies on the chemical essentials to the
discoloration of fish oil
J. Tokyo Univ. Fish. 43:127
- 670 Norris, F.A., J.P. Kass, and G.O. Burr. 1941
A study of dieneometry
Oil Soap 18:29-30
- 671 Nottevarp, O. 1961
Recent findings in fatty acid composition of
marine oils
In, Fish as Food, G. Borgstrom (Ed.), Academic
Press, N.Y.
- 672 Obst, Walter. 1936
Über die technische verwertung von vollfett
Dtsch. Apoth.-Ztg. 51:1584-1585
- 673 Ogata, K., S. Ohtake, H. Kawano, K. Kitagawa, H.
Tomimaga, and T. Akiyama. 1955
Experimental study on the toxicity of highly
unsaturated fatty acids.
Histopathologic reactions induced by this
material obtained from cuttlefish
Tokushima J. Exp. Med. 2:150-155
- 674 Okrugov, I.I., and S.D. Kopylenko. 1962
Arrangement for continuous hydrogenation of fats
U.S.S.R. Pat. 115,640 (addition to U.S.S.R. Pat.
103,217) issued Nov. 29, 1962
- 675 Olcott, H.S. 1959
The role of free fatty acids on antioxidant
effectiveness in unsaturated oils
J. Am. Oil Chem. Soc. 35:597-599
- * 676 Olcott, H.S. 1967
Antioxidants
In, Fish Oils, Stansby, M.E. (Ed.), AVI
Publishing Co., Westport, Conn. pp. 164-170
- 677 Olcott, H.S., and E. Einset. 1958
An antagonistic effect with antioxidants for
unsaturated fats
J. Am. Oil Chem. Soc. 35:159-160
- 678 Olcott, H.S., and H.A. Mattill. 1941
Constituents of fats and oils affecting the
development of rancidity
Chem. Rev. 29:257-437
- 679 Olcott, H.S., and J. Van der Veen. 1963
Role of individual phospholipids as antioxidants
J. Food Sci. 28:313-315
- 680 Oldfield, J.E., and A.P. Anglemeier. 1957
Feeding of crude and modified menhaden oil in
rations for swine
J. Anim. Sci. 16:917-921
- * 681 Oldfield, J.E., R.O. Sinnhuber, and A.A. Rasheed.
1963
Nutritive value of marine oils. II. Effects of
in vivo antioxidants in feeding menhaden oil to
swine
J. Am. Oil Chem. Soc. 40:357-360
- 682 Olley, J., and W.R.H. Duncan. 1965
Lipids and protein denaturation in fish muscle
J. Sci. Food Agric. 16:99-104
- 683 Olley, J., R. Pirie, and H. Watson. 1962
Lipase and phospholipase activity in fish
skeletal muscle and its relationship to protein
denaturation
J. Sci. Food Agric. 13:501-515
- 684 Olson, G., L.D. Matterson, and J. Tlustohowicz.
1962
Effect of vitamin E and antioxidants on fertility
and hatchability as affected by unsaturated
fatty acids
Poult. Sci. 41:675
- 685 Ong, I.T.L. 1969
Formation of aromatic compounds in hydrogenated
fish oils and their detection. (German)
Fette, Seifen, Antrichm. 71(12):1034-1038
- 686 Ono, T. 1939
Hydrolysis of fats and fat acid esters. III.
Hydrolysis of fats by enzymes
J. Agric. Chem. Soc. Jap. 15:1085-1096
- 687 Ozaki, J. 1926
Nutritive value of oils and fats. I
Bull. Agric. Chem. Soc. Jap. 2:10-31
- 688 Ozaki, J. 1926
Nutritive value of oils and fats. II
Bull. Agric. Chem. Soc. Jap. 2:845-865
- 689 Ozaki, J. 1927
Nutritive value of oils and fats. III
Bull. Agric. Chem. Soc. Jap. 3:977-1011
- 690 Ozaki, J. 1927
Nutritive value of oils and fats. IV
Bull. Agric. Chem. Soc. Jap. 3:1201-1205
- 691 Ozaki, J. 1932
Nutritive value of oils and fats. V
Bull. Agric. Chem. Soc. Jap. 8:1286-1303
- 692 Palfray, L. 1945
Hydrogenation of fatty substances. II.
Applications
Inds. Corps Gras 1:132-135
- * 693 Paluszak, J. 1967
Effect of extremely hydrogenated fat on the lipid
metabolism
Acta Physiol. Pol. 13(4):525-534
- 694 Panopoulos, G. 1932
Über eine spezifische reaktion der geharteten
fische
Praktika 7:325-326
- 695 Paschke, R.F., and D.H. Wheeler. 1955
Thermal polymerization of methyl linolenate,
alpha- and beta-eleostearate
J. Am. Oil Chem. Soc. 32:469-473
- 696 Pathak, S.P., and B.R. Reddy. 1964
The component acids of Indian "boundary" fish fats
J. Sci. Food Agric. 15:529-530

- 697 Pathak, S.P., and G.D. Pande. 1955
Component fatty acids of Indian shark liver oil
J. Am. Oil Chem. Soc. 32:7-9
- 8 Pathak, S.P., and P.M. Suwal. 1954
Component fatty acids of marine fish liver oils
J. Am. Oil Chem. Soc. 31:332-334
- 699 Patton, S., and G.W. Kurtz. 1955
2-thiobarbituric acid test for milk lipid oxidation
J. Dairy Sci. 38:901
- 700 Paul, S., and A. Roylance. 1962
Keeping properties of edible oils. Part I. The use of accelerated tests for assessment of the keeping properties of oils and the value of antioxidants
J. Am. Oil Chem. Soc. 39:163-165
- 701 Pedersen, T. 1938
On the course of hydrogenation in the industrial hardening of whale oil with nickel formate catalyst
Hvalradets Skrifter Det Norske Videnskaps-Akad. Oslo. 16:31
- 702 Peifer, J.J. 1966
Hypocholesterolemic effects induced in the rat by specific types of fatty acid unsaturation
J. Nutr. 88:351-388
- 703 Peifer, J.J. 1967
Hypercholesterolemic effects of marine oils
In, Fish Oils, M.E. Stansby (Ed.), AVI, Westport, Conn. pp. 322-361
- 704 Peifer, J.J., and R.T. Holman. 1955
Essential fatty acids, diabetes, and cholesterol
Arch. Biochem. Biophys. 57:520-521
- 705 Peifer, J.J., and R.T. Holman. 1959
Effect of saturated fat upon essential fatty acid metabolism in the rat
J. Nutr. 68:155-168
- 706 Peifer, J.J., and R.T. Holman. 1960
Acceleration of essential fatty acid deficiency by dietary cholesterol
J. Nutr. 70:411-417
- 707 Peifer, J.J., and W.O. Lundberg. 1959
Effect of unsaturated acids and fish oils on plasma and tissue lipids from hypercholesterolemic rats
Fed. Proc. 18:300
- 708 Peifer, J.J., P. Janssen, P. Ahn, W. Cox, and W.O. Lundberg. 1960
Studies on the distribution of lipids in hypercholesterolemic rats. 1. The effects of feeding palmitate, oleate, linoleate, linolenate, menhaden, and tuna oils
Arch. Biochem. Biophys. 86:302-308
- 709 Peifer, J.J., P. Janssen, R. Muesing, and W.O. Lundberg. 1962
The lipid depressant activities of whole fish and their component oils
J. Am. Oil Chem. Soc. 39:292-296
- 710 Peifer, J.J., W.O. Lundberg, S. Ishio, and E. Warmanen. 1965
Studies on the distribution of lipids in hypercholesterolemic rats. 3. Changes in hypercholesterolemia and tissue fatty acids induced by dietary fats and marine oil fractions
Arch. Biochem. Biophys. 110:270-283
- 711 Pelshenke, P., and A. Zeisset. 1937
Backeignungsprüfung einiger pflanzenfette und -ole und gehärteter waltrane
Mehl Brot 37(1):1-3
- 712 Peltzer, Josef. 1934
Vereinfachtes vakuumesterverfahren zum nachweis von margarine und gehärteten fetten
Z. Unters. Lebensm. 67:529-534
- 713 Peters, W.A., Jr. 1922
The efficiency and capacity of fractionating columns
Ind. Eng. Chem. 14:476-479
- 714 Peters, W.A., Jr. 1923
Calculations on fractionating columns from theoretical data
Ind. Eng. Chem. 15:402-403
- 715 Pigott, G.M. 1967
Production of fish oil
In, Fish Oils, M.E. Stansby (Ed.), AVI, Westport, Conn. pp. 183-192
- 716 Pihl, A. 1952
Cholesterol Studies. I. The cholesterol content of foods
Scand. J. Clin. Lab. Invest. 4:115
- 717 Pihl, A. 1955
On the mode of action of fatty acids in cholesterol absorption. Cholesterol studies: VI
Acta Physiol. Scand. 34:197-205
- * 718 Pohle, W.D., and R.L. Gregory. 1967
Standardization of nuclear magnetic resonance measurement of solids in fats and shortening
J. Am. Oil Chem. Soc. 44:397-399
- 719 Pohle, W.D., R.L. Gregory, and J.R. Taylor. 1962
A comparison of several techniques for prediction of relative stability of fats and oils to oxidation
J. Am. Oil Chem. Soc. 39:226-229
- 720 Pokorny, J., E. Mares, I. Zeman, and M. Karvanek. 1963
Methodes d'evaluation des matieres grasses hydrogenees
Potravin. Technol.
- 721 Pool, M.F., and A.A. Klose. 1951
Estimation of monocarbonyl compounds in rancid foods
J. Am. Oil Chem. Soc. 28:215
- 722 Post, O., and L.C. Craig. 1963
A new type of stepwise countercurrent distribution train
Anal. Chem. 35:641-647
- 723 Pringuet, A. 1964
Upgrading of fish oils
Oleagineux 19:411-416
- 724 Privett, O.S. 1957
Determination of structure and analysis of highly unsaturated and saturated acids of fish oils
Annu. Rep. Hormel Inst., Univ. of Minnesota, for 1955-1956, (Hormel Institute, Austin, Minn.) pp. 59-61
- 725 Privett, O.S., and E.C. Nickell. 1963
Preparation of highly purified fatty acids via liquid-liquid partition chromatography
J. Am. Oil Chem. Soc. 40:189-193
- 726 Privett, O.S., and M.L. Blank. 1964
Studies on the metabolism of linoleic acid in the essential fatty acid-deficient rat
J. Am. Oil Chem. Soc. 41:292-297
- 727 Privett, O.S., C. Nickell, W.O. Lundberg, and P.D. Boyer. 1955
Products of the lipoxidase-catalyzed oxidation of sodium linoleate
J. Am. Oil Chem. Soc. 32:505-511
- * 728 Privett, O.S., E. Aes-Jorgensen, R.T. Holman, and W.O. Lundberg. 1959
The effect of concentrates of polyunsaturated acids from tuna oil upon essential fatty acid deficiency
J. Nutr. 67:423-432

- 729 Privett, O.S., F.J. Pusch, and R.T. Holman. 1955
Polyethenoic fatty acid metabolism. VIII.
Nonpotency of cis-9, trans-12-linoleate as
essential fatty acid
Arch. Biochem. Biophys. 57:156
- 730 Privett, O.S., F.J. Pusch, T.R. Holman, and W.O.
Lundberg. 1960
Essential fatty acid properties of tuna, herring
and menhaden oils
J. Nutr. 71:66-69
- 731 Proctor, B.E., S.A. Goldblith, J.T.R. Nickerson,
and D.F. Parkas. 1960
Evaluation of the technical, economic, and
practical feasibility of radiation preservation
of fish
Report to U.S. Atomic Energy Commission
M.I.T. Contract No. AT(30-1)-2329
- 732 Pryde, E.H. 1951
Alcohols from menhaden oil by sodium reduction
J. Am. Oil Chem. Soc. 28:16-20
- 733 Quaife, M.L., N.W. Scrimshaw, and O.H. Lowry.
1949
A micromethod for assay of total tocopherols in
blood serum
J. Biol. Chem. 180:1229
- 734 Quin, L.P. 1965
The presence of compounds with carbon-phosphorus
bonds in some marine invertebrates
Biochem. 4:324-330
- 735 Quinby, G.E. (Ed.). 1972
Polychlorobiphenyls (PCB's) and related
chlorophenyls: effects on environment
Oak Ridge National Laboratory, Oak Ridge,
Tennessee (Oak Ridge Order No. TIRC-1, ORNL-EIS-
72-20) 1: bibliography 1881-1971
- 736 Radin, N.S., A.K. Hajra, and Y. Akahori. 1960
Preparation of methyl esters
J. Lipid Res. 1:250-251
- 737 Ralston, A.W. 1948
Fatty acids and their derivatives
John Wiley and Sons, N.Y.
- 738 Rapport, M.M., and N.F. Alonzo. 1960
The structure of plasmalogens. V. Lipids of
marine invertebrates
J. Biol. Chem. 235:1953-1956
- * 739 Rasheed, A.A., J.E. Oldfield, J. Kaufman, and
R.O. Sinnhuber. 1963
Nutritive value of marine oils I. Menhaden oil at
varying oxidation levels, with and without
antioxidants in rat diets
J. Nutr. 79:323-332
- 740 Raubut, M.M., H.A. Currier, A.M. Samsel, and V.P.
Wystrach. 1961
The free radical addition of phosphines to
unsaturated compounds
J. Org. Chem. 26:5138-5145
- 741 Raulin, J., and J. Petit. 1962
Nutritional value and physiopathological effect
of polymerized herring oil. I. Differences
between heating in air and heating in inert medium
Arch. Sci. Physiol. 16:77-87
- 742 Raulin, J., C. Richier, L. Escribano, and R.
Jacquet. 1959
Repercussions nutritionnelles et pathologiques de
l'usage alimentaire de l'huile de poisson
desodorisée par chauffage. Description d'une
myocardite infiltrante
Compt. Rend. Acad. Sci. (Paris) 248:1229
- 743 Raulin, J., C. Richier, and R. Jacquet. 1962
Nutritional and pathological effects of
nutritional use of fish oils deodorized by heat
Symp. Substances Etrangères Aliments 5, Budapest,
1959 pp. 227-233
- 744 Reed, S.A. 1964
Some nutritional aspects of cod liver oil. I.
Its essential fatty acid
and hypocholesterolaemic activity
J. Sci. Food Agric. 15:399-407
- 745 Reed, S.A. 1965
Fish oils, their composition, processing, and
some industrial applications
J. Oil Colour Chem. Assoc. 48:213-214
- * 746 Reichsman, P. No date
Survival time of hypertensive rats receiving fish-
oil extracts
Science 104(2690):64
- 747 Reinert, R.E. 1969
Insecticides and the Great Lakes
Lirnos Magazine (U.S.) 2(3):4-9
- 748 Reiser, R., B. Stevenson, M. Kayama, R.B.R.
Choudhury, and D.W. Hood. 1963
The influence of dietary fatty acids and
environmental temperature on the fatty
acid composition of teleost fish
J. Am. Oil Chem. Soc. 40:507-513
- 749 Renninger, George. 1958
Effect of hydrogenated fish oils on citrus-tree
destroying nematodes
Plant Dis. Rep. 42(9):1057-1065
- 750 Reppe, W., and H. Kroper. 1953
Carbonylation. II. Carboxylic acids and their
derivatives from olefinic compounds and carbon
monoxide
Liebig's Ann. 582:38-71
- 751 Rini, J. 1960
Refining, bleaching, stabilization deodorization,
and plasticization of fats, oils and shortening
J. Am. Oil Chem. Soc. 37:512-520
- 752 Robinson, K.L., and W.E. Coey. 1951
A brown discoloration of pig fat and vitamin E
deficiency
Nature, London 168:997-998
- 753 Rodegker, W. and J.C. Nevenzel. 1964
The fatty acid composition of three marine
invertebrates
Comp. Biochem. Physiol. 11:53-60
- 754 Roe, E.T., and D. Swern. 1960
Branched carboxylic acids from long-chain
unsaturated compounds and carbon monoxide at
atmospheric pressure
J. Am. Oil Chem. Soc. 37:661-668
- 755 Roe, E.T., G.R. Riser, and D. Swern. 1961
Preparation, characterization and evaluation of
mono- and diesters of carboxystearic acid
J. Am. Oil Chem. Soc. 38:527-531
- 756 Rose, A. 1936
Distillation efficiency in 3- and 6-mm
fractionating columns
Ind. Eng. Chem., Ind. Ed. 28:1210-1212
- 757 Rosenthal, M.L. 1964
Biological role and practical uses of squalene
and squalane
In, Cosmetics and the Skin, Lubow and Wells
(Eds.), Reinhold, N.Y. pp. 646-652
- 758 Rost, E., G. Sonntag, and A. Weitzel. 1937
Zur gesundheitlichen beurteilung einiger in der
neuzeit fur genusszwecke empfohlener fette. II.
Stoffwechselversuche mit geharteten tierischen
fett (waltran), sowie mit freien und mit an
athylalkohol (athylester) und an glykol
(glykolester) gebundenen
hochmolekularen fettsauren (stearin-, palmitin-,
olsaure)
Arch. Hyg. Bakt. 118:139-205

- 759 Roubal, W.T. 1963
Tuna fatty acids: II. Investigations of the composition of raw and processed domestic tuna
J. Am. Oil Chem. Soc. 40:215-218
- 60 Roubal, W.T., and A.L. Tappel. 1966
Damage to proteins, enzymes and amino acids by peroxidizing lipids
Arch. Biochem. Biophys. 113(58):5-8
- 761 Rouser, G., G. Kritchevsky, D. Heller, and E. Lieber. 1963
Lipid composition of beef brain, beef liver, and sea anemone: Two approaches to quantitative fractionation of complex lipid mixtures
J. Am. Oil Chem. Soc. 40:425-454
- 762 Rowan, M.K. 1958
Chloromyxum in juvenile pilchards
Fish. Ind. Res. Inst., Annu. Rep. 12:10-11
- 763 Saba, J.G., M.A. Nielsen, and A.K. Sumner. 1970
Effect of commercial processing techniques on lindane- and DDT-C-14 residues in rapeseed oil
J. Agric. Food Chem. 18(1):43-44
- 764 Saiki, M., S. Pang and T. Mori. 1959
Studies on the lipid of plankton-II. Fatty acid composition of lipids from euphausiacea collected in the Antarctic and northern Pacific oceans
Bull. Jap. Soc. Sci. Fish. 24:837-839
- 765 Sakurai, Y., Y. Kato, T. Masuhara, and S. Hayakawa 1954
On nutritive value of finwhale oil and polymerized oil. I
J. Jap. Soc. Food Nutr. 3:155-156
- 766 Sanchez-Torres, J.R. 1967
Fish oil industry in South America
In, Fish oils, M.E. Stansby (Ed.), AVI, Westport, Conn. pp. 394-404
- 767 Sand, G. 1965
Herring oil and seal industry
Tidsskr. Kjem. Bergves. Metall. 5:150-152
- 768 Sand, G. 1967
Fish oil industry in Europe
In, Fish Oils, M.E. Stansby (Ed.), AVI, Westport, Conn. pp. 405-421
- 769 Sanford, F. Bruce, and C.F. Lee. 1961
Menhaden industry
U.S. Fish and Wildlife Service, Commercial Fish. TL 31:31 pp.
- 770 Sanford, F.B., and C.F. Lee. 1960
U.S. Fish reduction industry
Technical Leaflet, Bureau of Commercial Fisheries, Washington, D.C. No. 14
- 771 Schilling, K. 1957
Isolierung und Bestimmung partiell hydrierte Fettsauren mittels der Quecksilber-Adducte
Pette-Seife-Anstrichm. 61:765
- 772 Schlenk, H. 1961
Crystallization of fatty acids
J. Am. Oil Chem. Soc. 38:728-736
- 773 Schlenk, H., and D.M. Sand. 1962
Collection of gas-liquid chromatography fractions by gradient cooling
Anal. Chem. 34:1676
- 774 Schlenk, H., and J.L. Gellerman. 1960
Esterification of fatty acids with diazomethane on a small scale
Anal. Chem. 32:1412-1414
- 775 Schlenk, H., and J.L. Gellerman. 1961
Column chromatography of fatty acids
J. Am. Oil Chem. Soc. 38:555-562
- 776 Schlenk, H., and M.A. Ener. 1959
Solubility and fractionation of lipids in sulfur dioxide
J. Am. Oil Chem. Soc. 36:145-149
- 777 Schlenk, H., and R.T. Holman. 1953
Methanolysis of triglycerides by an anion exchange resin
J. Am. Oil Chem. Soc. 30:103-104
- 778 Schlenk, H., J.L. Gellerman, and D.M. Sand. 1962
Acylation cyclodextrins as stationary phases for comparative gas-liquid chromatography
Anal. Chem. 34:1529-1532
- 779 Schlenk, H., J.L. Gellerman, J.A. Tillotson, and H.K. Mangold. 1957
Paper chromatography of lipids
J. Am. Oil Chem. Soc. 34:377
- 780 Schlenk, H., W. Sen, and D.M. Sand. 1963
Interconversion of long-chain fatty acids in the rat
Biochim. Biophys. Acta 70:708-710
- 781 Schmid, H.H.O., H.L. Mangold, and W.O. Lundberg. 1965
Critical solution temperatures of aliphatic compounds. I. Homologous and vinylogous fatty acids and methyl esters
Microchem. J. 9:134-144
- 782 Schmidt, E.G. 1930
The phenylation of oleic acid
J. Am. Chem. Soc. 52:1172-1174
- 783 Scholfield, C.R. 1961
Countercurrent distribution
J. Am. Oil Chem. Soc. 38:562-565
- 784 Scholfield, C.R., E.P. Jones, and H.J. Dutton. 1961
Hydrogenation of linolenate
Anal. Chem. 33:1745
- 785 Scholfield, C.R., E.P. Jones, J. Nowakowska, E. Selke, and H.J. Dutton. 1961
Hydrogenation of linolenate. II. Hydrazine reduction
J. Am. Oil Chem. Soc. 38:208-211
- 786 Scholfield, C.R., E.P. Jones, R.O. Butterfield, and H.J. Dutton. 1963
Hydrogenation of linolenate
Anal. Chem. 35:386
- 787 Scholfield, C.R., J. Nowakowska, and H.J. Dutton. 1960
Preparation of pure fatty acid methyl esters by countercurrent distribution
J. Am. Oil Chem. Soc. 37:27-30
- 788 Schreiber, M.L., G.E. Vail, R.M. Conrad, and L.F. Payne. 1947
Effect of tissue-fat stability on deterioration of frozen poultry
Poult. Sci. 26:14-19
- 789 Schultz, H.W., E.A. Day, and R.O. Sinnhuber. 1962
Symposium on foods: lipids and their oxidation
AVI Publishing Co., Westport, Conn.
- 790 Scott, M.L. 1951
Studies on the enlarged hock disorder in turkeys. 3. Evidence of the detrimental effect of fish liver oil and the beneficial effect of dried brewers' yeast and other materials
Poult. Sci. 30:846-855
- 791 Scott, M.L. 1953
Prevention of enlarged hock disorder in turkeys with niacin and vitamin E
Poult. Sci. 32:670-677
- 792 Scott, P.P. and E.R. Humphreys. 1961
Addition of herring and vegetable oils to diet of cats
Proc. Nutr. Soc. 21(1):xiii

- 793 Sen, N., and H. Schlenk. 1964
The structure of polyenoic odd-and even-numbered fatty acids of mullet (*Mugil cephalus*)
J. Am. Oil Chem. Soc. 41:241-247
- 94 Senior, J.R. 1964
Intestinal absorption of fats
J. Lipid Res. 5:495-521
- 795 Sergeev, A.G. 1956
Installation for continuous hydrogenation of fats
U.S.S.R. Pat. 103,217 issued Jul. 25, 1956
- 796 Shewan, J.M. 1962
The influence of irradiation preservation on the nutritive value of fish and fishery products
In, Fish in Nutrition, E. Heen and R. Kreuzer (Eds.), Fishing News (Books) Ltd., London
- 797 Shimamura, U. 1964
Effect of addition of dimethylpolysiloxane on hydrogenation of marine animal fatty oils
J. Jap. Oil Chem. Soc. 13:255-259
- 798 Shimizu, M. 1951
Lipolytic activity of pancreatic lipase on dietary fats
J. Jap. Biochem. Soc. 23:103-105
- 799 Shima, Y., and H. Taguchi. 1964
A comparative study of fatty acid composition of fish
Bull. Jap. Soc. Sci. Fish. 30:179-188
- 800 Shima, Y., and H. Taguchi. 1964
A comparative study of fatty acid compositions of wild and cultivated "ayu" sweet smelt (*Plecoglossus altivelis*)
Bull. Jap. Soc. Sci. Fish. 30:918-925
- 801 Shuster, C.Y., J.R. Froines, and H.S. Olcott. 1964
Phospholipids of tuna white muscle
J. Am. Oil Chem. Soc. 41:36-41
- 802 Siakotos, A.N., and G. Rouser. 1965
Analytical separation of nonlipid water soluble substances and gangliosides from other lipids by dextran gel column chromatography
J. Am. Oil Chem. Soc. 42:913-919
- 803 Silk, H.M., H.H. Sephton, and H.H. Hahn. 1954
South African pilchard oil. 2. Concentrates of highly unsaturated fatty acids and alcohols derived from South African pilchard oil
Biochem. J. 57:574-577
- 804 Silk, M.H., and A.J. DeKoning. 1964
Phospholipids of South African pilchard (*Sardinia ocellata jenyns*)
J. Am. Oil Chem. Soc. 41:619-622
- 805 Silk, M.H., and H.H. Hahn. 1954
South African pilchard oil. 4. The isolation and structure of a hexadecatetraenoic acid from South African pilchard oil
Biochem. J. 57:582-587
- 806 Sinclair, G.D. 1964
Fatty acid sizing of hardboards
Tappi 47:579-583
- 807 Sinclair, R.G. 1932
The metabolism of the phospholipids. 5. The relationship between the amount of fat ingested and the degree of unsaturation of the phospholipids and neutral fat in the tissues of the rat
J. Biol. Chem. 96:103-125
- 808 Singleton, W.S. 1960
Solution properties
In, Fatty Acids, 2nd Ed., K.S. Markley (Ed.), Interscience Publishers, N.Y. Part I
- 809 Singsen, E.P., L.D. Matterson, A. Kozeff, R.H. Bunnell, and E.L. Jungherr. 1954
Studies on encephalomalacia in the chick. I. The influence of a vitamin E deficiency on the performance of breeding hens and their chicks
Poult. Sci. 33:192-201
- 810 Singsen, E.P., L.M. Potter, L.D. Matterson, R.H. Bunnell, A. Kozeff, and E.L. Jungherr. 1954
Studies on encephalomalacia in the chick. 4. The influence of oil in fish meal and oils from various species of fish on incidence of encephalomalacia
Poult. Sci. 33:1081
- 811 Singsen, E.P., L.M. Potter, L.D. Matterson, R.H. Bunnell, A. Kozeff, and E.L. Jungherr. 1955
Studies on encephalomalacia in the chick. 4. Influence of oil in fish meal, oils from various species of fish, and animal fats on the incidence of encephalomalacia
Poult. Sci. 34:1075-1079
- 812 Singsen, E.P., R.H. Bunnell, L.D. Matterson, A. Kozeff, and E.L. Jungherr. 1955
Studies on encephalomalacia in the chick. 2. The protective action of diphenyl-p-phenylenediamine against encephalomalacia
Poult. Sci. 34:262-271
- 813 Sinnhuber, R.O., and T.C. Yu. 1958
2-Thiobarbituric acid method for the measurement of rancidity in fishery products. II. The quantitative determination of malonaldehyde
Food Technol. 12:9-12
- 814 Sinnhuber, R.O., T.C. Yu, and T.C. Yu. 1958
Characterization of the red pigment formed in the 2-thiobarbituric acid determination of oxidative rancidity
Food Res. 23:626
- 815 Skau, E.L., and R.E. Boucher. 1954
An interpolative method of calculating solubilities of missing members of homologous series
J. Phys. Chem. 58:460-468
- * 816 Smith, K.J., P.B. Polen, D.M. DeVries, and F.B. Coon. 1968
Removal of chlorinated pesticides from crude vegetable oils by simulated commercial processing procedures
J. Am. Oil Chem. Soc. 45:866-869
- 817 Snyder, F. No date
The biochemistry of lipids containing an ether bond
Progress in Chemistry of Fats and Other Lipids, R.T. Holmar, Ed.
- 818 Sokoloff, B., M. Toyomizu, W. Trauner, and G. Renninger. 1959
The fungicidal activity of the unsaturated fatty acids and quaternary salts prepared from fish oils
J. Am. Oil Chem. Soc. 36:234-237
- 819 Sone, H. 1967
Fish oil industry in Japan
In, Fish Oils, M.E. Stansby (Ed.), AVI, Westport, Conn. pp. 422-427
- 820 Sonntag, N.O.V. 1961
Halogenation, dehalogenation, and dehydrohalogenation
In, Fatty Acids, Their Chemistry, Properties, Production, and Uses, 2nd Ed., K.S. Marley (Ed.), Interscience Publishers, N.Y.
- 821 Spanhoff, R.W., and H.C. Jacobsen. 1950
Bacterial action of cod-liver oil in the treatment of wounds
Pharm. Weekbl. 85:742-750
- 822 Sperry, W.M., and M. Webb. 1950
A revision of the Schoenheimer-Sperry method for cholesterol determination
J. Biol. Chem. 187:97
- 823 Sreenivasan, B., and J.B. Brown. 1956
Isomerization of linoleic, linolenic, and other polyunsaturated acids with potassium tertiary butoxide and its application in the spectrophotometric estimation of these acids
J. Am. Oil Chem. Soc. 33:521-526

- 824 Stacy, F.W., and J.E. Harris, Jr. 1963
Formation of carbon-hetero atom bonds by free radical chain additions to carbon-carbon multiple bonds
In, *Org. Reactions*, John Wiley and Sons, N.Y. 43:150-376
- 825 Stander, G.H. 1956
Size vs. fat content of pilchards
Fish. Ind. Res. Inst. Annu. Rep. 10:16
- 826 Stansby, M.E. (Ed.). 1963
Industrial Fishery Technology
The Reinhold Publishing Company, Inc., N.Y. 393 pp.
- 827 Stansby, M.E. (Ed.). 1967
Fish oils, their chemistry, technology, stability, nutritional properties, and uses
The AVI Publishing Company, Inc., Westport Conn. 440 pp.
- * 828 Stansby, M.E. Accepted for publication 1973
Problems discouraging use of fish oil in American manufactured shortening and margarine
J. Am. Oil Chem. Soc.
- 829 Stansby, M.E. 1935
Analysis of fish. I. Tests for the condition of the oil of fish flesh
J. Assoc. Offic. Agric. Chem. 18:646-621
- 830 Stansby, M.E. 1962
Speculations on "fishy" odors
Food Technol. 16(4):28-32
- 831 Stansby, M.E. 1962
Proximate composition of fish
In, *Fish in Nutrition*, Heen and Kreuzers (Eds.), Fishing News, London pp. 55-60
- * 832 Stansby, M.E. 1967
Misconceptions about nutritional properties of fish oils
In, *Fish Oils*, Stansby, M.E. (Ed.), AVI Publishing Co., Westport, Conn. pp. 283-288
- * 833 Stansby, M.E. 1967
Appendix-physical and chemical constants of fish oils
In, *Fish Oils*, Stansby, M.E. (Ed.), AVI Publishing Co., Westport, Conn. pp. 433-434
- 834 Stansby, M.E. 1967
Odors and flavors
In, *Fish Oils*, M.E. Stansby (Ed.), AVI, Westport, Conn. pp. 174-180
- * 835 Stansby, M.E. 1967
General analytical methods
In, *Fish Oils*, Stansby, M.E. (Ed.), AVI Publishing Co., Westport, Conn. pp. 43-51
- * 836 Stansby, M.E. 1969
Nutritional properties of fish oils
World Review of Nutrition and Dietetics 11:46-105
- 837 Stansby, M.E., and A.S. Hall. 1967
Chemical composition of commercially important fish of the United States
Fish. Ind. Res. 3(4):29-46
- 838 Stansby, M.E., and G. Jellinek. 1965
Flavor and odor characteristics of fishery products with particular reference to early oxidative changes in menhaden oil
In, *The Technology of Fish Utilization*, R. Kreuzer, (Ed.), The Fishing News (Books) Ltd., London pp. 171-178
- * 839 Stansby, M.E., and G.M. Knobl, Jr. 1968
U.S. Fish meal and oil industry
Draft
- 840 Stansby, M.E., P.E. Zollman, and R.K. Winkelmann. 1967
Efficacy of fish oils in healing wounds and burns
Fish. Ind. Res. 3(4):25-27
- 841 Steenbock, H., M.H. Irwin, and J. Weber. 1936
The comparative rate of absorption of different fats
J. Nutr. 12:103-111
- 842 Steinberg, G., W.H. Slaton, Jr., and D.R. Howton. 1957
Metabolism of essential fatty acids. V. Metabolic pathway of linolenic acid
J. Biol. Chem. 224:841-849
- 843 Stirton, A.J., B.B. Schaeffer, A.A. Stawitzke, J.K. Weil, and W.C. Ault. 1948
Aryl acids from oleic acid. Variables affecting the yield and properties
J. Am. Oil Chem. Soc. 25:365-368
- 844 Stoffel, W. 1963
Der Stoffwechsel der ungesättigten Fettsäuren. I. Zur Biosynthese hochungesättigter Fettsäuren
Z. Physiol. Chem. 333:71-88
- 845 Stoffel, W., and E.H. Ahrens, Jr. 1958
Isolation and structure of the C16 unsaturated fatty acids in menhaden body oil
J. Am. Chem. Soc. 80:6604-6608
- 846 Stoffel, W., and F.H. Ahrens, Jr. 1960
The unsaturated fatty acids in menhaden body oil: the C18, C20, and C22 series
J. Lipid Res. 1:139-146
- 847 Stoffel, W., F. Chu, and E.H. Ahrens, Jr. 1959
Analysis of long-chain fatty acids by gas-liquid chromatography. Micro method for preparation of methyl esters
Anal. Chem. 31:307-308
- * 848 Stout, V.F. 1968
Pesticide levels in fish of the northeast Pacific
Bull. Environ. Contam. Toxicol. 3(4):240-246
- * 849 Stout, V.F., F.L. Beezhold, and C.R. Houle. 1970
DDT residue levels in some U.S. fishery products and the effectiveness of some treatments in reducing them
FAO Technical Conference on Marine Pollution and its Effects on Living Resources and Fishing
Rome, Italy, Dec. 9-18, 1970
- 850 Strohecker, R., R. Vaubel, and A. Tenner. 1937
Die bestimmung des aktiven sauerstoffs und die anwendung dieser methode bei der untersuchung von fetten
Fette Seifen 44:246-250
- 851 Sumerwell, W.N. 1957
Liquid-solid countercurrent distribution of fatty acids with urea
J. Am. Chem. Soc. 79:3414-3415
- 852 Suzuki, K. 1960
Preparation of a coating material by chlorination of fish oil. VII. Preparation of a chlorination product with high melting point
Bull. Jap. Soc. Sci. Fish. 26:509-513
- 853 Suzuki, K. 1965
The pattern of mammalian brain gangliosides. III. Regional and developmental differences
J. Neurochem. 12:969-979
- 854 Svennerholm, L. 1964
The gangliosides
J. Lipid Res. 5:445-155
- 855 Swain, L.A. 1953
Fatty acid composition of fish oils. II. Herring oil
Fish. Res. Board Can. Progr. Rep. Pac. Coast Sta. 94:24-26
- 856 Swain, L.A. 1954
Fatty acid composition of fish oils. III. Sockeye salmon offal oil
Fish. Res. Board Can., Progr. Rep. Pac. Coast Sta. 99:6-8

- 857 Swain, L.A. 1958
The composition and properties of fish oils
In, Progress in the Chemistry of Fats and other
Lipids, R.T. Holman, W.O. Lundberg and T. Malkin
(Eds.), Pergamon Press, London
- 858 Swell, L., H. Field, Jr., and C.R. Treadwell.
1960
Correlation of arachidonic acid of serum
cholesterol esters in different species
with susceptibility to atherosclerosis
Proc. Soc. Exp. Biol. Med. 104:325-328
- 859 Swell, L., M.D. Law, P.E. Schools, Jr., and C.R.
Treadwell. 1961
Tissue lipid fatty acid changes following the
feeding of high carbohydrate, essential fatty
acid-supplemental diets to rabbits
J. Nutr. 75:181-191
- 860 Swell, L., P.E. Schools, Jr., and C.R. Treadwell.
1962
Correlation of diet and serum cholesterol level
with serum lipid fatty acid composition
Proc. Soc. Exp. Biol. Med. 109:682-685
- 861 Swern, D. 1961
Chemical oxidations
In, Fatty Acids, Their Chemistry, Properties,
Production, and Uses, 2nd Ed., K.S. Markley (Ed.)
Interscience Publishers, N.Y.
- 862 Swift & Co. 1962
Epoxy monoglycerides and their esters
Brit. Pat. 908,500 issued Oct. 17, 1962
- 863 Takama, K., M. Hatano, K. Zama, and H. Igarashi.
1963
Gas chromatographic analysis of fatty acid
composition of marine animal oils
Bull. Faculty Fish. Hokkaido Univ. 14:31-36
- 864 Takano, M. and Y. Kikuya. 1938
Studies on the partial hydrogenation of fish oil.
IX. The hydrogenation of herring oils
J. Soc. Chem. Ind., Jap. 41(7):238B-240B
- 865 Takano, Masakichi. 1935
Studien über die partielle Hydrierung von
fischölen. III. Hydrierung der
ungesättigten fettsäuren der ölsäurereihe des
sardinenöles
J. Soc. Chem. Ind., Jap. (Suppl.) 38:302B-304B
- 866 Takano, Masakichi. 1935
Studien über die partielle Hydrierung von
fischölen. V. Bildung von
ungesättigter fettsäuren der linolensäurereihe
bei der Hydrierung von methylklupanodonat
J. Soc. Chem. Ind., Jap. (Suppl.) 38:652B-653B
- 867 Takano, Masakichi, and Yoshio Kumeno. 1935
Studien über die partielle Hydrierung von
fischölen. II. Der allgemeine verlauf
der Hydrierung von sardinenöl
J. Soc. Chem. Ind., Jap. (Suppl.) 38:252B-254B
- 868 Takemura, K.H., and L.A. Goldblatt. 1957
The migration of double bonds during the
isomerization of methyl linoleate with palladium
on carbon catalyst
J. Am. Oil Chem. Soc. 34:15-17
- 869 Tappel, A.L. 1955
Studies of the mechanism of vitamin E action. 3.
In vitro copolymerization of oxidized fats with
protein
Arch. Biochem. 50:266-280
- 870 Tarladgis, B.G., and B.M. Watts. 1960
Malonaldehyde production during the controlled
oxidation of pure unsaturated fatty acids
J. Am. Oil Chem. Soc. 37:403
- 871 Tchetcheroff, Elisee. 1935
Das problem der identifizierung und bestimmung
des milchfettes
Congr. Chim. Ind. Brux. 15(1):396-407
- 872 Teeter, H.M. 1948
Bromination of methyl esters of soybean fat acids
and methyl linoleate with N-bromosuccinimide
J. Am. Oil Chem. Soc. 25:243-245
- 873 Teeter, H.M., J.L. O'Donnell, W.J. Schneider,
L.E. Gast, and M.J. Danzig. 1957
Reactions of conjugated fatty acids. IV. Diels-
Alder adducts of 9,11-octadecadienoic acid
J. Org. Chem. 22:512-514
- 874 Thomasson, H.J. 1953
Biological standardization of essential fatty
acids (a new method)
Int. Rev. Vitam. Res. 35:62
- 875 Thomasson, H.J. 1953
Biological standardization of essential fatty
acids (a new method)
Int. Rev. Vitam. Res. 25:62
- 876 Thomasson, H.J. 1953
Results obtained by a new biological method of
standardizing essential fatty acids
In, International Conference on Biochemical
Problems of Lipides, Palais der Academien,
Brussels p. 212
- 877 Thomasson, H.J. 1956
Biological value of oils and fats. IV. Rate of
intestinal absorption
J. Nutr. 59:343-352
- 878 Thomasson, H.J. 1956
Biological value of various natural oils and fats
In, Biochemical Problems of Lipids, Popjack and
Le Breton (Eds.), Interscience Publishers, N.Y.
- 879 Thomasson, H.J. 1962
Essential fatty acids
Nature 194:973
- 880 Thompson, M.H. 1964
Cholesterol content of various species shellfish.
I. Method of analysis and preliminary survey
variables
Fish. Ind. Res. 2(3):11
- 881 Tidwell, H.C., and L.E. Holt. 1936
The estimation of the total lipids and the lipid
partition in feces
J. Biol. Chem. 112:605
- 882 Tidwell, H.C., J.C. McPherson, and W.W. Burr, Jr.
1962
Effect of the saturation of fats upon the
disposition of ingested cholesterol
Am. J. Clin. Nutr. 11:108-114
- 883 Tobolsky, A.V., D.J. Metz, and R.B. Mesrobian.
1950
Low temperature autoxidation of hydrocarbons:
the phenomenon of maximum rates
J. Am. Chem. Soc. 72:1942-1952
- 884 Toyama, K. 1962
Relation between the efficiency of antioxidant
and the degree of unsaturation of substrate oil
J. Tokio Univ. Fish. 48:111-118
- 885 Toyama, T., and T. Kaneda. 1962
Nutritive aspects of fish oils
In, Fish as Food, G. Borgstrom (Ed.), Academic
Press, London Vol. 2
- 886 Toyama, Y. and T. Takagi. 1954
Fatty oils of aquatic invertebrates. IV. Fatty
acids and sterols in the fat of Spisula
sachalinensis
Nippon Kagaku Zasshi 75:1238-1241
- 887 Toyama, Y., and T. Takagi. 1962
Eicosapentaenoic and docosahexaenoic acid in the
cuttlefish oil
Fette-Seifen-Anstrichm. 64:134-137

- 888 Toyama, Y., and T. Tsuchiya. 1925
Separation of highly unsaturated fatty acids from fish oil
J. Soc. Chem. Ind. Jap. 28:963-965
- 889 Toyomizu, M., K. Kawasaki, and Y. Tomiyasu. 1963
Effect of dietary oil on the fatty acid composition of rainbow trout oil
Bull. Jap. Soc. Sci. Fish. 29:957-961
- 890 Tsuchiya, T. 1961
Biochemistry of fish oils
In, Fish as Food, G. Borgstrom (Ed.), Academic Press, N.Y.
- 891 Tsuchiya, Tomotaro, and Goroku Akiyama. 1933
Umesterung von fetten mit glycerin
J. Soc. Chem. Ind., Jap. (Suppl.) 36:233B-234B
- 892 Tsuchiya, Y., and M. Kayama. 1957
Studies on the conjugated fatty acids. Part II. The thermal polymerization of the conjugated fish oil obtained by the nickel-on-carbon catalyst isomerization
Tohoku J. Agric. Res. 7:277-289
- 893 Tuna, N., and H.K. Mangold. 1964
Fatty acids of the atherosclerotic plaque
In, Evolution of the Atherosclerotic Plaque, R.L. Jones (Ed.), Univ. of Chicago Press
- 894 Turner, W.A., E.B. Meigs, and H.T. Converse. 1936
Toxic effect of cod-liver oil in the ration of the rabbit and the calf
J. Biol. Chem. 114:civ
- 895 Turpeinen, O.J. 1938
Further studies on the unsaturated fatty acids essential in nutrition
J. Nutr. 15:351-366
- * 896 U.S. Bureau of the Census, Department of Commerce. 1965, 1969, 1971
U.S. Imports for Consumption and General Imports
Government Printing Office, Washington, D.C.
Publications PT 246-71, 246-69, 246-65
- * 897 U.S. Department of Agriculture. 1972
Agricultural Statistics
U.S. Government Printing Office, Washington, D.C.
- * 898 U.S. Pharmacopoeial Commission. 1960
U.S. Pharmacopoeia, 18th Edition
U.S. Pharmacopoeial Commission, Bethesda, Maryland
- * 899 U.S. Tariff Commission. No date
Synthetic Organic Chemicals, U.S. Production and Sales
Government Printing Office, Washington, D.C.
- * 900 Ueno, S. and S. Nakaguchi. 1936
On the cooling curves of oils, fats and waxes in their natural states as well as hardened and paraffins. I. (Fr.)
J. Soc. Chem. Ind., Jap. 39:79B-82B
- * 901 Ueno, S. 1938
The relationship between thiocyanogen value and iodine value in the course of oil-hardening
J. Soc. Chem. Ind., Jap. 41(6):200B-202B
- * 902 Ueno, S., M. Yamashita and Y. Ota. 1928
On the nutritive value of hardened oils. II
J. Soc. Chem. Ind., Jap. 31:92-93
- 903 Ueno, S., T. Hikada, and T. Okamoto. 1955
The occurrence of highly unsaturated, triethanoid, and diethanoid alcohols in the liver oils of *Laemonema morosum*
J. Jap. Oil Chem. Soc. 4:26-27
- 904 Ueno, Sei-ichi and Kiyoji Yasuhara. 1962
Beschaffenheit der japanischen fischole vom standpunkt der hartung
J. Soc. Chem. Ind., Japan 30:388-389
- 905 Ueno, Sei-ichi and Riichiro Yamasaki. 1932
Über die neuen, bei der hydrierung von fischolen gebildeten verbindungen. IV. Die Fettsäuren
J. Soc. Chem. Ind., Jap. (Suppl.) 35:492B-493
- 906 Ueno, Sei-ichi. 1930
Über die neuen bei der hydrierung von fischolen entstehenden kohlenwasserstoffe
J. Soc. Chem. Ind., Jap. (Suppl.) 33:264B-266B
- 907 Ueno, Sei-ichi, and Riichiro Yamasaki. 1930
Über neue bei der hydrierung von fischolen gebildete verbindungen. II. Kohlenwasserstoffe
J. Soc. Chem. Ind., Jap. (Suppl.) 33:451B-452B
- 908 Ueno, Sei-ichi, and Sumio Matsuda. 1935
Über die fraktionierte destillation der fettsäuren von extrem hochgeharteten olen. II
J. Soc. Chem. Ind., Jap. (Suppl.) 38:691B-692B
- 909 Underhill, F.P., and O.E. Closson. 1905
The physiological behavior of methylene blue and methylene azure: a contribution to the study of the oxidation and reduction processes in the animal organism
Am. J. Physiol. 13:358-371
- 910 Urdahl, N., H. Saav, and A. Helgerud. 1972
Hydrogenation of fish oils produced from raw materials at different levels of freshness
Paper presented at Eleventh World Congress, ISF, Goteborg, Sweden, June, 1972
- 911 Vagelos, P.R. 1964
Lipid metabolism
Ann. Rev. Biochem. 33:139-172
- 912 Van Der Merwe, R.P., and L. Kapp. 1961
Oxidation phenomena in the oil of South African fish meal
Fish. Ind. Res. Inst., Cape Town, S. Afr. Annu. Rep. pp. 56-57
- * 913 Van Der Steur, J.P.K. 1961
Nutritional aspects of processing and chemical additives on fats and oils
Fed. Proc. 20(Suppl. 7):217-223
- 914 Vandenheuvel, F.A., and D.R. Vatcher. 1956
Partition chromatography of aliphatic acids: Quantitative resolution of normal chain even acids from C12 to C24
Anal. Chem. 28:838-845
- 915 Vendell, J.H., and J.W. Putnam. 1945
Fish oil flavor in eggs
Poult. Sci. 24:285-286
- 916 Vilbushevitch, M. 1912
Converting fat, fish oil, etc. into compounds of higher melting point by hydrogen and a catalyzer
U.S. Pat. 1,020,758 issued Apr. 30, 1912
- 917 Von Rudloff, E. 1956
Periodate-permanganate oxidation. IV. Determination of the position of double bonds in unsaturated fatty acids and esters
J. Am. Oil Chem. Soc. 33:126
- 918 Wakil, S.J. 1961
Mechanism of fatty acid synthesis
J. Lipid Res. 2:1-24
- 919 Wallace, J.C., W.R. Peterson, A.F. Chadwick, and D.O. Barlow. 1958
Hydroxylation of methyl oleate. A new, direct method
J. Am. Oil Chem. Soc. 35:205-207
- 920 Walling, C., and E.S. Huyser. 1963
Free radical additions to olefins to form carbon-carbon bonds
In, Org. Reactions, John Wiley and Sons, N.Y. 13:91-149
- 921 Ward, R.J. 1963
Sex differences in the alpha-tocopherol requirement of rats as shown by the hemolysis test
Brit. J. Nutr. 17:135

- 922 Ward, R.J., and C.C. Amos. 1962
Growth of rats given cod liver oil
Brit. J. Nutr. 16:219-222
- 923 Watanabe, Y., Y. Ito, and Matsuura. 1957
Oxidation of unsaturated fatty acids. I.
Reaction of methyl oleate with selenium dioxide
J. Sci. Hiroshima Univ., Ser. A 20:203-208
- 924 Waterman, H.I., C. Boelhouwer, and L.J. Revallier. 1951
Hydrogenation of fatty oils
Elsevier Publishing Co., N.Y. p. 241
- 925 Weitkamp, A.W. 1947
The amplified distillation of methyl esters of fatty acids
J. Am. Oil Chem. Soc. 24:236-238
- 926 Weitkamp, A.W. 1955
Distillation
J. Am. Oil Chem. Soc. 32:640-646
- 927 Wheeler, D.H. 1932
Peroxide formation as a measure of auto-oxidative deterioration
Oil Soap 9:89-97
- 928 Whitcutt, J.M. 1954
The chemistry of pilchard oils
S. Afr. Ind. Chem. 11(9):175-176
- 929 Whitcutt, J.M. 1957
South African pilchard oil. 6. The isolation and structure of a docosahexaenoic acid from South African pilchard oil
Biochem. J. 67:60-64
- 930 Whitcutt, J.M., and D.A. Sutton. 1956
South African pilchard oil. 5. The isolation and structure of an eicosapentaenoic acid from South African pilchard oil
Biochem. J. 63:469-475
- 931 White, H.B. Jr., and W.L. Williams. 1965
Analysis of unsaturated fatty esters as mercurated adducts by thin-layer chromatography
Fed. Proc. 24:662
- 932 White, H.B., Jr., and F.W. Quackenbush. 1959
A simplified technique for analysis by alkali isomerization
J. Am. Oil Chem. Soc. 36:653-656
- 933 White, H.B., Jr., and F.W. Quackenbush. 1962
Isolation of pure linolenate as its mercuric acetate adduct
J. Am. Oil Chem. Soc. 39:547-549
- 934 White, H.B., Jr., and F.W. Quackenbush. 1962
Separation of fatty estermercuric acetate adducts on alumina
J. Am. Oil Chem. Soc. 39:511-513
- 935 Whitehouse, M.W. 1964
Drugs, hormones, and other factors influencing steroid and sterol metabolism
In, Lipid Pharmacology, R. Paoletti, (Ed.), Academic Press, N.Y.
- 936 Will, P. 1961
New solvent system for separation of fatty acids C10-C18 by countercurrent distribution
Anal. Chem. 33:647-648
- 937 Willard, J.G., and M.L. Martinez. 1961
The selective hydrogenation of linolenic acid
J. Am. Oil Chem. Soc. 38:282-286
- 938 Williamson, B., and L.C. Craig. 1947
Identification of small amounts of organic compounds by distribution studies. Calculation of theoretical curves
J. Biol. Chem. 168:687-697
- 939 Winstein, S., and H.J. Lucas. 1938
The coordination of silver ion with unsaturated compounds
J. Am. Chem. Soc. 60:836-847
- 940 Wisniak, J., A. Cancino, and J.C. Vega. 1964
Epoxidation of anchovy oils. A study of variables
Ind. Eng. Chem., Prod. Res. Dev. 3:304-311
- 941 Wolfe, D.A., R. Venkata, and D.G. Cornwell. 1965
Studies on the fatty acid composition of crayfish lipids
J. Am. Oil Chem. Soc. 42:633-637
- 942 Wolff, J. 1921
Über einige tranhartfette
Z. Dtsch. Öl- Fettind. 41:179-180
- 943 Wolfrom, M.L., and G.P. Arsenault. 1960
Chromatographic separation of 2,4-dinitrophenylhydrazine derivatives of highly oxygenated carbonyl compounds
Anal. Chem. 32:693
- 944 Wood, J.D., and J. Biely. 1960
The effect of dietary marine fish oils on the serum cholesterol levels in hypercholesterolemic chickens
Can. J. Biochem. Physiol. 38:19-24
- 945 Woodford, P.P., and G.M. Van Gent. 1960
Gas-liquid chromatography of fatty acid methyl esters: the carbon numbers as a parameter for comparison of columns
J. Lipid Res. 1:188-190
- 946 Worne, H.E., and L.W. Smith. 1959
Effects of certain pure long-chain polyunsaturated fatty acid esters on the blood lipids of man
Am. J. Med. Sci. 237:710-721
- 947 Wurster, C.F., Jr., J.H. Copenhaver, Jr., and P.E. Shafer. 1963
Separation of the methyl esters of oleic, linoleic, and linolenic acids by column chromatography using cation exchange resin containing silver ion
J. Am. Oil Chem. Soc. 40:513-514
- * 948 Wyatt, C.J., and E.A. Day. 1963
Autoxidation of fish oils. II. Changes in the carbonyl distribution of autoxidizing salmon oils
J. Food Sci. 28:305-312
- 949 Yamada, M. 1961
Studies on the lipid of plankton. I. Lipid of Neomysis nakazawai
J. Jap. Oil Chem. Soc. 10:236-239
- 950 Yendo, Yoshiomi. 1935
Über den Mechanismus der Kohlenwasserstoffbildung durch Hydrierung von Fettsäuren bei hoher Temperatur und hohem Druck
Sci. Pap. Inst. Phys. Chem. Res. 28(603/606)
- 951 Young, R.J., and R.L. Garret. 1963
Effect of oleic and linoleic acids on the absorption of fatty acids in the chick
J. Nutr. 81:321-329
- 952 Yu, T.C., and R.O. Sinnhuber. 1957
2-Thiobarbituric acid method for the measurement of rancidity in fishery products
Food Technol. 11:104-108
- 953 Yu, T.C., E.A. Day, and R.O. Sinnhuber. 1961
Autoxidation of fish oils. I. Identification of volatile monocarbonyl compounds from auto-oxidizing salmon oil
J. Food Sci. 26:192-197
- 954 Yudkin, J. 1964
Dietary fat and dietary sugar in relation to ischaemic heart disease and serum lipids
Lancet 2:4-5
- 955 Zama, K. 1964
Phospholipids of aquatic animals. A symposium on marine lipid metabolism
Bull. Jap. Soc. Sci. Fish. 30:660-672

956 Zuckerman, L., and G.H. Marquardt. 1963
Muscle, erythrocyte and plasma electrolytes and
some other muscle constituents of rabbits with
nutritional muscular dystrophy
Proc. Soc. Exp. Biol. Med. 112:609

57 Zunini, Enrico. 1933
Beitrag zur untersuchung von butter mittels einer
thermometrischen prufung der loslichkeit der
fettsauren ihrer bestandteile
Ann. Chim. Appl. 23:557-567